

Color path-integral Monte-Carlo simulations of quark-gluon plasma: Thermodynamic and transport properties

V.S. Filinov,^{1,*} Yu.B. Ivanov,² V.E. Fortov,¹ M. Bonitz,³ and P.R. Levashov¹

¹*Joint Institute for High Temperatures, Russian Academy of Sciences, Moscow, Russia*

²*Kurchatov Institute, Kurchatov sq. 1, Moscow, Russia*

³*Institute for Theoretical Physics and Astrophysics, Christian Albrechts University, Kiel, Germany*

Based on the quasiparticle model of the quark-gluon plasma (QGP), a color quantum path-integral Monte-Carlo (PIMC) method for calculation of thermodynamic properties and – closely related to the latter – a Wigner dynamics method for calculation of transport properties of the QGP are formulated. The QGP partition function is presented in the form of a color path integral with a new relativistic measure instead of the Gaussian one traditionally used in the Feynman-Wiener path integral. A procedure of sampling color variables according to the SU(3) group Haar measure is developed for integration over the color variable. It is shown that the PIMC method is able to reproduce the lattice QCD equation of state at zero baryon chemical potential at realistic model parameters (i.e. quasiparticle masses and coupling constant) and also yields valuable insight into the internal structure of the QGP. Our results indicate that the QGP reveals *quantum liquid-like* (rather than gas-like) properties up to the highest considered temperature of 525 MeV. The pair distribution functions clearly reflect the existence of gluon-gluon bound states, i.e. glueballs, at temperatures just above the phase transition, while meson-like $q\bar{q}$ bound states are not found. The calculated self-diffusion coefficient agrees well with some estimates of the heavy-quark diffusion constant available from recent lattice data and also with an analysis of heavy-quark quenching in experiments on ultrarelativistic heavy ion collisions, however, appreciably exceeds other estimates. The lattice and heavy-quark-quenching results on the heavy-quark diffusion are still rather diverse. The obtained results for the shear viscosity are in the range of those deduced from an analysis of the experimental elliptic flow in ultrarelativistic heavy ions collisions, i.e. in terms the viscosity-to-entropy ratio, $1/4\pi \lesssim \eta/S < 2.5/4\pi$, in the temperature range from 170 to 440 MeV.

PACS numbers: 12.38Mh, 31.15.Qg, 51.20.+d, 52.27Gr

Keywords: quark-gluon plasma, diffusion, viscosity, equation of state, path-integral Monte-Carlo simulations

I. INTRODUCTION

Determining the properties of the quark-gluon plasma (QGP) is nowadays one of the most important goals in high-energy nuclear physics. In recent years, experiments at the Relativistic Heavy-Ion Collider (RHIC) at Brookhaven National Laboratory [1] and the Large Hadron Collider (LHC) at CERN [2] have provided a wealth of data from which one can obtain information on a number of features of the QGP. The most striking result, obtained from analysis of these experimental data [1, 3], is that the deconfined quark-gluon matter behaves as an almost perfect fluid rather than a perfect gas, as it could be expected from the asymptotic freedom.

There are various approaches for a theoretical study of the QGP each of which has its advantages and disadvantages. The most fundamental way to compute the properties of strongly interacting matter is provided by the lattice QCD [4–6]. In particular, the lattice QCD computations [7] indicated that the transition into the QGP phase at small chemical potentials has a nature of a rapid crossover, which, strictly speaking, is not a phase transition. However, following the custom of the heavy-ion society, we call this transformation as a phase transition of the crossover type. Interpretation of these very complicated computations requires application of various QCD motivated, albeit schematic, models simulating various aspects of the full theory. Moreover, such models are needed in cases when the lattice QCD fails, e.g. at large quark chemical potentials and out of equilibrium. While certain progress has been achieved in recent years, transport properties of the QGP are still poorly accessible within lattice QCD – only viscosity for pure gluonic matter was computed [8]. It is, therefore, crucial to devise reliable and manageable theoretical tools for a quantitative description of non-Abelian QGP both in and out of equilibrium.

A semi-classical approximation, based on a quasiparticle picture has been introduced in Refs. [9–13]. It is motivated by the expectation that the main features of non-Abelian plasmas can be understood in simple semi-classical terms

*Corresponding author E-mail: vladimir_filinov@mail.ru

without the difficulties inherent to a full quantum field-theoretical analysis. Independently the same ideas were implemented in terms of molecular dynamics (MD) [14]; this approach was further developed in a series of works [15, 16]. The MD allows one to treat soft processes in the QGP which are not accessible by perturbative means.

Strongly correlated behavior of the QGP is expected to show up in long-ranged spatial correlations of quarks and gluons which, in fact, may give rise to liquid-like and, possibly, solid-like structures. This expectation is based on a very similar behavior observed in electrodynamic plasmas [15, 17, 18]. This similarity was exploited to formulate a classical nonrelativistic model of a color Coulomb interacting QGP [15] which was numerically analyzed by classical MD simulations. Quantum effects were either neglected or incorporated phenomenologically via a short-range repulsive correction to the pair potential. Such a rough model may, however, become a critical issue at high densities. Similar models in electrodynamic plasmas showed poor behavior in the region of strong wave function overlap, in particular around the Mott density where bound states break up. For temperatures and densities of the QGP considered in Ref. [15] these effects are very important since the quasiparticle thermal wave length is of the order of the average interparticle distance. Therefore, to account for quantum effects, we follow an idea of Kelbg [19] that allows one to rigorously include quantum corrections to the pair potential¹. Strictly speaking, this method is applicable only to weak coupling. To extend the method to stronger couplings, an “improved Kelbg potential” was derived, which contains a single free parameter, being fitted to the exact solution of the quantum-mechanical two-body problem. Using the method of the improved Kelbg potential in classical MD simulations one is able to describe thermodynamic properties of a partially ionized plasma up to moderate couplings [21]. However, this approach may fail, if bound states of more than two particles are formed in the system. This is a result of break-down of the pair approximation for the density matrix, as demonstrated in Refs. [21]. A superior approach, which does not have this limitation, consists in the use of the original Kelbg potential in path integral Monte Carlo simulations (PIMC) which effectively map the problem onto a high-temperature weakly coupled and weakly degenerate one. This allows one to advance the analysis to strong couplings and is, therefore, a relevant choice for the present purpose. The PIMC method has been successfully applied to various phases of strongly coupled electrodynamic plasmas [22, 23]. Examples are partially-ionized dense hydrogen plasmas, where liquid-like and crystalline behavior was observed [24–26], as well as electron-hole plasmas in semiconductors [27, 28], including excitonic bound states.

In this paper we extend the previous classical nonrelativistic simulations [15] in two ways: first, we include quantum and spin effects and, second, we take into account the dominant relativistic effects, cf. section II. This is done in the frame of quantum Monte Carlo simulations where we rewrite the partition function of this system in the form of color path integrals with a new relativistic measure instead of Gaussian one used in Feynman-Wiener path integrals. For the integration of the partition function over color variables we develop a procedure of sampling the color quasiparticle variables according to the SU(3) group Haar measure with the quadratic and cubic Casimir conditions. The developed approach self-consistently takes into account the Fermi (Bose) statistics of quarks (gluons). The main goal of this article is to test the present Color Path-Integral Monte-Carlo (PIMC) against known lattice data [4] and to predict additional properties of the QGP, which are still inaccessible from lattice QCD. First results of the path integral approach for thermodynamic properties of the nonideal QGP have been already briefly reported in Ref. [29] for SU(3) group and in Refs. [30–33] for SU(2) group. In this paper we show that the PIMC method is able to reproduce the QCD lattice equation of state at vanishing baryon-charge density and also yields valuable insight into the internal structure of the QGP. These results are presented in section IV A.

Hydrodynamic simulations of relativistic heavy-ion collisions require not only knowledge of thermodynamic properties of the QGP but also of the transport properties. Unfortunately the PIMC method itself is not able to directly predict transport properties. Therefore, to simulate quantum QGP transport and thermodynamic properties within a unified approach we combine the path integral and Wigner (in phase space) formulations of quantum mechanics (section III). There the kinetic coefficients are calculated by means of Kubo formulas. In this approach the PIMC method is used to generate initial conditions (equilibrium quasiparticle configurations) for dynamical trajectories describing the time evolution for spatial, momentum and color variables. Correlation functions and kinetic coefficients are calculated as averages of Weyl’s symbols of dynamic operators along these trajectories. The basic ideas of this approach have been published in Ref. [34]. Using this approach we calculate the self-diffusion coefficient and viscosity of the strongly coupled QGP. These results are presented in section IV B.

¹ The idea to use a Kelbg-type effective potential also for quark matter was independently proposed by K. Dusling and C. Young [20]. However, their potentials are limited to weakly nonideal systems.

II. THERMODYNAMICS OF QGP

In this section we summarize the main ideas of our approach to the thermodynamic properties of the strongly correlated quark-gluon plasma. This approach is based on a generalization of the Feynman path integral representation of quantum mechanics to high energy matter. Before deriving the main equations of our PIMC approach we specify the simplifications and model parameters.

A. Basics of the model

The basic assumptions of the model are similar to those of Ref. [15]:

- I. Quasiparticles masses (m) are of order or higher than the mean kinetic energy per particle. This assumption is based on the analysis of QCD lattice data [35–37]. For instance, at zero net-baryon density it amounts to $m \gtrsim T$, where T is a temperature.
- II. In view of the first assumption, interparticle interaction is dominated by a color-electric Coulomb potential. Magnetic effects are neglected as subleading ones.
- III. Relying on the fact that the color representations are large, the color operators are substituted by their average values, i.e. by Wong's classical color vectors (8D in $SU(3)$) with the quadratic and cubic Casimir conditions [38].
- IV. We consider the 3-flavor quark model. Since the masses of the 'up', 'down' and 'strange' quarks extracted from lattice data are still indistinguishable, we assume these masses to be equal. As for the gluon quasiparticles, we allow their mass to be different (heavier) from that of quarks.

The quality of these approximations and their limitations were discussed in Ref. [15]. Thus, this model requires the following quantities as a function of temperature (T) and quark chemical potential (μ_q) as an input:

- 1. quasiparticle masses, for quarks m_q and gluons m_g , and
- 2. the coupling constant g^2 , or $\alpha_s = g^2/4\pi$.

Input quantities should be deduced from lattice QCD data or from an appropriate model simulating these data.

It has been established that hard modes (in terms of *hard thermal loop* approximation [39–41]) behave like quasiparticles [41]. Therefore, masses of these quasiparticles should be deduced from nonperturbative calculations taking into account hard field modes, e.g., they can be associated with pole masses deduced from lattice QCD calculations. At the same time, the soft quantum fields are characterized by very high occupation numbers per mode. Therefore, to leading order, they can be well approximated by soft classical fields. This is precisely the picture we are going to utilize: massive quantum quasiparticles (hard modes) interacting via classical color fields. Applicability of such approach was discussed in Refs. [10, 15] in detail. Our approach differs from that of Ref. [10, 15] by a quantum treatment to quasiparticles instead of the classical one, and additionally by a relativistic description of the kinetic energy instead of the nonrelativistic approximation of Ref. [10].

B. Color Path Integrals

We consider a multi-component QGP consisting of N color quasiparticles: N_g gluons, N_q quarks and \overline{N}_q antiquarks. The Hamiltonian of this system is $\hat{H} = \hat{K} + \hat{U}^C$ with the kinetic and color Coulomb interaction parts

$$\hat{K} = \sum_i \sqrt{\hat{\mathbf{p}}_i^2 + m_i^2(T, \mu_q)}, \quad \hat{U}^C = \frac{1}{2} \sum_{i \neq j} \frac{g^2(T, \mu_q)(Q_i \cdot Q_j)}{4\pi|\mathbf{r}_i - \mathbf{r}_j|}. \quad (1)$$

Here i and j summations run over quark and gluon quasiparticles, $i, j = 1, \dots, N$, $N = N_q + \overline{N}_q + N_g$, $N_q = N_u + N_d + N_s$ and $\overline{N}_q = \overline{N}_u + \overline{N}_d + \overline{N}_s$ are total numbers of quarks and antiquarks of all flavours (*up*, *down* and *strange*), respectively, 3D vectors \mathbf{r}_i are quasiparticle spatial coordinates, the Q_i denote the Wong's quasiparticle color variable (8D-vector in the group $SU(3)$), $(Q_i \cdot Q_j)$ denote scalar product of color vectors. Nonrelativistic approximation for potential energy is used, while for kinetic energy we still keep relativistic form as the quasiparticle masses are not negligible as compared with temperature. The eigenvalue equation of this Hamiltonian is usually called the spinless

Salpeter equation. It may be regarded as a well-defined approximation to the BetheSalpeter formalism [42] for the description of bound states within relativistic quantum field theories, obtained when assuming that all bound-state constituents interact instantaneously and propagate like free particles [43]. Among others, it yields semirelativistic descriptions of hadrons as bound states of quarks [44, 45].

In the classic approximation this system is governed by Wong's equations of motion [38]

$$\frac{d\mathbf{p}_i(t)}{dt} = \mathbf{F}_i(t), \quad (2)$$

$$\frac{d\mathbf{r}_i(t)}{dt} = \mathbf{v}_i(t), \quad (3)$$

$$\frac{dQ_i^a(t)}{dt} = \Upsilon_i^a(t), \quad (4)$$

where \mathbf{p}_i is the momentum of a quasiparticle, $\mathbf{v}_i = \mathbf{p}_i / \sqrt{\mathbf{p}_i^2 + m_i^2}$ is its velocity, $\mathbf{F}_i = -\partial U^C / \partial \mathbf{r}_i$ is the color-electric force experienced by the quasiparticle, and

$$\Upsilon_i^a = \sum_{j \neq i} \sum_{b,c} \check{f}_{abc} \frac{g^2 Q_i^b Q_j^c}{4\pi |\mathbf{r}_i - \mathbf{r}_j|}, \quad (5)$$

is the driving force in equation of motion for the color charge, \check{f}_{abc} are structure constants of the group SU(3) and $a, b, c = 1, \dots, 8$ (see Appendix I). Below we consider spatial degrees of freedom quantum-mechanically while the color dynamics, still classically.

Thermodynamic properties in the grand canonical ensemble with given temperature (T), net-quark-number (μ_q) and strange (μ_s) chemical potentials, and fixed volume V are fully described by the grand partition function²

$$Z(\mu_q, \mu_s, \beta, V) = \sum_{\{N\}} \frac{\exp\{\mu_q(N_q - \bar{N}_q)/T\} \exp\{\mu_s(N_s - \bar{N}_s)/T\}}{N_u! N_d! N_s! \bar{N}_u! \bar{N}_d! \bar{N}_s! N_g!} Z(\{N\}, V, \beta), \quad (6)$$

$$Z(\{N\}, V, \beta) = \sum_{\sigma} \int_V dr d\mu Q \rho(r, Q, \sigma; \{N\}; \beta), \quad (7)$$

where $\{N\} = \{N_u, N_d, N_s, \bar{N}_u, \bar{N}_d, \bar{N}_s, N_g\}$, $\rho(r, Q, \sigma; \{N\}; \beta)$ denotes the diagonal matrix elements of the density-matrix operator $\hat{\rho} = \exp(-\beta \hat{H})$ with $\beta = 1/T$. Here r , σ and Q denote the multi-dimensional vectors related spatial, spin and color degrees of freedom, respectively, of all quarks, antiquarks and gluons. The σ summation and spacial ($dr \equiv d^3r_1 \dots d^3r_N$) and color ($d\mu Q \equiv d\mu Q_1 \dots d\mu Q_N$) integrations run over all individual degrees of freedom of the particles, $d\mu Q_i$ denotes integration over SU(3) group Haar measure, see Appendix I. Usual choice of the strange chemical potential is $\mu_s = -\mu_q$ (nonstrange matter), such that the total factor in front of $(N_s - \bar{N}_s)$ is zero. Therefore, below we omit μ_s from the list of variables.

Since the masses and the coupling constant depend on the temperature and quark chemical potential, special care should be taken to preserve thermodynamical consistency of this approach. To achieve this, thermodynamic functions such as pressure, P , entropy, S , baryon number, N_B , and internal energy, E , should be calculated through respective derivatives of the logarithm of the partition function

$$P = \partial(T \ln Z) / \partial V, \quad (8)$$

$$S = \partial(T \ln Z) / \partial T, \quad (9)$$

$$N_B = (1/3) \partial(T \ln Z) / \partial \mu_q, \quad (10)$$

$$E = -PV + TS + 3\mu_q N_B. \quad (11)$$

This is a conventional way of maintaining the thermodynamical consistency in approaches of the Ginzburg–Landau type as they are applied in high-energy physics, e.g., in the PNJL model.

² Here we explicitly write sum over different quark flavors (u,d,s). Below the sum over quark degrees of freedom is understood in the same way.

The exact density matrix $\rho = e^{-\beta\hat{H}}$ of interacting quantum systems can be constructed using a path integral approach [46, 47] based on the operator identity $e^{-\beta\hat{H}} = e^{-\Delta\beta\hat{H}} \cdot e^{-\Delta\beta\hat{H}} \dots e^{-\Delta\beta\hat{H}}$, where the r.h.s. contains $n+1$ identical factors with $\Delta\beta = \beta/(n+1)$, which allows us to rewrite³ the integral in Eq. (7) in the form

$$\begin{aligned}
& \sum_{\sigma} \int dr^{(0)} d\mu Q \rho(r^{(0)}, Q, \sigma; \{N\}; \beta) \\
&= \sum_{\sigma} \int d\mu Q dr^{(0)} dr^{(1)} \dots dr^{(n+1)} \rho^{(1)} \rho^{(2)} \dots \rho^{(n)} \\
&\times \sum_{P_q} \sum_{P_{\bar{q}}} \sum_{P_g} (-1)^{\kappa_{P_q} + \kappa_{P_{\bar{q}}}} \mathcal{S}(\sigma^{(0)}, \hat{P}_q \hat{P}_{\bar{q}} \hat{P}_g \sigma^{(n+1)}) \hat{P}_q \hat{P}_{\bar{q}} \hat{P}_g \rho^{(n+1)} \Big|_{\sigma^{(n+1)} = \sigma^{(0)}} \delta(r^{(n+1)} - r^{(0)}) \\
&\equiv \int d\mu Q \int dr^{(0)} dr^{(1)} \dots dr^{(n+1)} R(r^{(0)}, r^{(1)}, \dots, r^{(n+1)}; Q; \{N\}; \beta) \delta(r^{(n+1)} - r^{(0)}). \tag{12}
\end{aligned}$$

Notice that the color charge Q is a classical variable already in the mixed (i.e. coordinate-momentum) representation, see Appendix I. Therefore, we do not build a chain of n different Q -variables. The spin variable σ is the same in all $\rho^{(l)}$ except for the $\rho^{(n+1)}$, where it is initially set $\sigma^{(n+1)}$ and only after permutations performed is put to σ . The spin gives rise to the spin part of the density matrix (\mathcal{S}) with exchange effects accounted for by the permutation operators \hat{P}_q , $\hat{P}_{\bar{q}}$ and \hat{P}_g acting on spatial $r^{(n+1)}$ degrees of freedom and the spin projections $\sigma^{(n+1)}$. The sum runs over all permutations with parity κ_{P_q} and $\kappa_{P_{\bar{q}}}$. In Eq. (12)

$$\rho^{(l)} = \rho(r^{(l-1)}, r^{(l)}, Q; \{N\}; \Delta\beta) = \langle r^{(l-1)}, Q | e^{-\Delta\beta\hat{H}} | r^{(l)}, Q \rangle, \tag{13}$$

is an off-diagonal element of the density matrix. Accordingly each quasiparticle is represented by a set of coordinates $\{r_i^{(0)}, \dots, r_i^{(n)}\}$ ("beads") and a 8-dimensional color vector Q_i in the $SU(3)$ group. Thus, all "beads" of each quasiparticle are characterized by the same spin projection, flavor and color charge. Notice that masses and coupling constant in each $\rho^{(l)}$ are the same as those for the original quasiparticles, i.e. these are still defined by the actual temperature T .

The main advantage of decomposition (12) is that it allows us to use perturbation theory to obtain approximation for density matrices $\rho^{(l)}$, which is applicable due to smallness of artificially introduced factor $1/(n+1)$. This means that in each $\rho^{(l)}$ the ratio $g^2(T, \mu_q)(Q_i \cdot Q_j)/[4\pi|r_i^{(l)} - r_j^{(l)}|T(n+1)]$ can be always made much smaller than one, which allows us to use perturbation theory with respect to the potential. Each factor in Eq. (12) should be calculated with the accuracy of order of $1/(n+1)^\theta$ with $\theta > 1$, as in this case the error of the whole product in the limit of large n will be equal to zero. In the limit $(n+1) \rightarrow \infty$, ρ^l can be approximated by a product of two-particle density matrices $\rho_{ij}^{(l)}$ [23, 46, 47]. This approximation can be deduced from operator expansion

$$\exp(-\Delta\beta\hat{H}) = \exp(-\Delta\beta\hat{K}) \exp(-\Delta\beta\hat{U}^C) \exp(-\Delta\beta^2 [\hat{K}, \hat{U}^C]/2) \dots \tag{14}$$

As the first approximation with the error proportional to $(1/(n+1))^2$, we can write

$$\exp(-\Delta\beta\hat{H}) \approx \exp(-\Delta\beta\hat{K}) \exp(-\Delta\beta\hat{U}^C), \tag{15}$$

It is very important that in this case the error of the whole product in Eq. (12) is proportional to the $1/(n+1)$ and is equal to zero in the limit $n \rightarrow \infty$. The second advantage of the decomposition in Eq. (12) is that it reduces quantum multi-particle interaction to the pair-wise sum of two-particle interactions described by two-particle classical density matrices in each $\rho^{(l)}$. Thus, neglecting the commutator terms in Eq. (14), we arrive at the following expression for the density matrix of Eq. (13)

$$\rho_{\text{neglecting commutators}}^{(l)} = \rho_0^{(l)} \exp[-\Delta\beta U^C(r^{(l)}, Q)], \tag{16}$$

where $\rho_0^{(l)}$ is the corresponding density matrix of noninteracting particles. This approximation works well for potentials bounded below. However, the Coulomb potential can go to minus infinity and hence the result (16) diverges in this limit.

³ For the sake of notation convenience, we ascribe superscript (0) to the original variables.

A more sophisticated treatment is required to avoid this divergence. All the calculations along this line can be rigorously performed for the two-particle density matrix $\rho^{[2]}(r, r'; Q; \Delta\beta)$, where $r = \{\mathbf{r}_1, \mathbf{r}_2\}$, $r' = \{\mathbf{r}'_1, \mathbf{r}'_2\}$ and $Q = \{Q_1, Q_2\}$. Expanding the two-particle density matrix up to the second order in $1/(n+1)$, one arrives to the following result [19]

$$\begin{aligned} \rho^{[2]}(r, r'; Q; \Delta\beta) &\approx \rho_0^{[2]}(r, r'; Q; \Delta\beta) \\ &- \int_0^1 d\tau \int d^3\tilde{\mathbf{r}} \frac{\Delta\beta g^2(Q_i \cdot Q_j)}{4\pi|\tilde{\mathbf{r}}|\Delta\lambda_{12}^2\sqrt{\tau(1-\tau)}} \exp\left(-\frac{\pi|\mathbf{r}_{12}-\tilde{\mathbf{r}}|^2}{\Delta\lambda_{12}^2(1-\tau)}\right) \exp\left(-\frac{\pi|\tilde{\mathbf{r}}-\mathbf{r}'_{12}|^2}{\Delta\lambda_{12}^2\tau}\right) \\ &\approx \rho_0^{[2]} \exp[-\Delta\beta \Phi_{12}(\mathbf{r}_{12}, \mathbf{r}'_{12}, Q_1, Q_2)], \end{aligned} \quad (17)$$

where $\mathbf{r}_{12} = \mathbf{r}_1 - \mathbf{r}_2$, $\mathbf{r}'_{12} = \mathbf{r}'_1 - \mathbf{r}'_2$, $\Delta\lambda_{12} = \sqrt{2\pi\Delta\beta/m_{12}}$ is defined in terms of the reduced mass of the pair of particles: $m_{12} = m_1 m_2 / (m_1 + m_2)$, and $\rho_0^{[2]}$ is the two-particle density matrix of noninteracting particles. In the end of Eq. (17) the result is presented in the form similar to Eq. (16), i.e. in terms of an off-diagonal two-particle effective quantum potential Φ_{12} , which is called a Kelbg potential [19]. Eq. (17) is the definition of the color Kelbg potential. The diagonal part of the color Kelbg potential can be obtained analytically

$$\Phi_{12}(\mathbf{r}_{12}, \mathbf{r}_{12}, Q_1, Q_2) \approx \frac{g^2(Q_i \cdot Q_j)}{4\pi\Delta\lambda_{12}x_{12}} \left\{ 1 - e^{-(x_{12})^2} + \sqrt{\pi}x_{12} [1 - \text{erf}(x_{12})] \right\}, \quad (18)$$

where $x_{12} = |\mathbf{r}_{12}|/\Delta\lambda_{12}$. Notice that the color Kelbg potential approaches the color Coulomb potential at distances larger than $\Delta\lambda_{12}$. What is of prime importance, the color Kelbg potential is finite at zero distance, thus removing in a natural way the classical divergences and making any artificial cut-offs, often applied (see, e.g., Ref. [15]), obsolete. This color potential is a straightforward generalization of the corresponding potential of electromagnetic Coulomb plasmas [21]. The off-diagonal color Kelbg potential can be approximated by the diagonal ones by means of $\Phi_{12}(\mathbf{r}_{12}, \mathbf{r}'_{12}, Q_1, Q_2) \approx [\Phi_{12}(\mathbf{r}_{12}, \mathbf{r}_{12}, Q_1, Q_2) + \Phi_{12}(\mathbf{r}'_{12}, \mathbf{r}'_{12}, Q_1, Q_2)]/2$.

Unfortunately such rigorous consideration of multiparticle density matrix for particles interacting by potentials unbounded below is not available. Therefore, following the experience gained in electromagnetic Coulomb plasmas, we use the following widely used ansatz [46, 47], which generalizes Eq. (17):

$$\rho^{(l)} = \rho_0^{(l)} \exp \left[-\Delta\beta \frac{1}{2} \sum_{i,j(i \neq j)}^N \Phi_{ij} \left(\mathbf{r}_i^{(l-1)} - \mathbf{r}_j^{(l-1)}, \mathbf{r}_i^{(l)} - \mathbf{r}_j^{(l)}, Q_i, Q_j \right) \right]. \quad (19)$$

Now we are able to construct R of Eq. (12). The density matrix of noninteracting particles should be expressed in terms of determinants and permanents of single-particle density matrices in the standard way. Generalizing the electrodynamic plasma results [23] to the quark-gluon plasma case, we write approximate R

$$\begin{aligned} R(r^{(0)}, r^{(1)}, \dots, r^{(n+1)}; Q; \{N\}; \beta) &= \\ &= \exp\{-\beta U\} \sum_{\sigma} \left[\prod_{l=1}^n \prod_{i=1}^N \phi_{ii}(\mathbf{r}_i^{(l-1)}, \mathbf{r}_i^{(l)}, \Delta\beta) \right] \\ &\times \frac{\text{per} \|\phi(r^{(n)}, r^{(0)}, \Delta\beta)\|_{N_g}}{\Lambda_g^{3(n+1)N_g}(\Delta\beta)} \frac{\det \|\phi(r^{(n)}, r^{(0)}, \Delta\beta)\|_{N_q}}{\Lambda_q^{3(n+1)N_q}(\Delta\beta)} \frac{\det \|\phi(r^{(n)}, r^{(0)}, \Delta\beta)\|_{\overline{N}_q}}{\Lambda_{\overline{q}}^{3(n+1)\overline{N}_q}(\Delta\beta)}, \end{aligned} \quad (20)$$

In Eq. (20) the effective total color interaction energy is

$$U = \frac{1}{n+1} \sum_{l=1}^{n+1} \frac{1}{2} \sum_{i,j(i \neq j)}^N \Phi_{ij} \left(\mathbf{r}_i^{(l-1)} - \mathbf{r}_j^{(l-1)}, \mathbf{r}_i^{(l)} - \mathbf{r}_j^{(l)}, Q_i, Q_j \right). \quad (21)$$

Other quantities in Eq. (20) are defined as follows:

$$\Lambda_a^3(\beta) = \lambda_a^3 \sqrt{\pi/2(\beta m_a)^5} \quad (22)$$

with $\lambda_a = \sqrt{2\pi\beta/m_a}$ being a thermal wavelength of an a type quasiparticle ($a = q, \overline{q}, g$). The antisymmetrization and symmetrization are taken into account by the symbols “det” and “per” denoting the determinant and permanent,

respectively. Eq. (20) is exact in the limit of $n \rightarrow \infty$. Indeed, since each factor in Eq. (12) has an error of order of $1/(n+1)^\theta$ with $\theta > 1$, the error of the whole product in the limit of $n \rightarrow \infty$ equals zero. Matrix $\phi(r, r', \Delta\beta)$ is defined by its matrix elements

$$\phi_{ij}(r, r', \Delta\beta) = \frac{K_2[z_{ij}(r, r', \Delta\beta)]}{[z_{ij}(r, r', \Delta\beta)]^2} \delta[\delta_{a_i, g} + (\delta_{a_i, q} + \delta_{a_i, \bar{q}}) \delta_{f_i, f_j} \delta_{\sigma_i, \sigma_j}] \quad (23)$$

with

$$z_{ij}(r, r', \Delta\beta) = \Delta\beta m_i \sqrt{1 + |\mathbf{r}_i - \mathbf{r}'_j|^2 / \Delta\beta^2} \quad (24)$$

defined in terms of the modified Bessel function K_2 . These matrix elements are nonzero only for particles of the same type, i.e. $a_i = a_j$. Additional Kronecker symbols in spin, σ_i , and flavor, f_i , indices of the particles are applicable only to quark and antiquark matrix elements. They prevent Pauli blocking for particles with different spins and flavors. The quantity ϕ describes the *relativistic measure* of trajectories in the color path integral. This measure is associated with relativistic operator of kinetic energy in Eq. (1). In the limit of large mass this measure coincides with the Gaussian one used in Feynman-Wiener path integrals. Due to factors $\delta_{a_i, a_j} \delta_{f_i, f_j}$ the matrix ϕ has a block structure corresponding to different types of particles and different flavors of quarks and antiquarks. Subscripts N_a near det and per operations precisely refer to the corresponding blocks, which in case of quarks and antiquarks are still subdivided into sub-blocks related to flavors.

The dominant contribution to the partition function comes from configurations in which the “size” of the quasiparticle cloud of ‘beads’ is of the order of the Compton wavelength $\lambda_C = 1/m_i$. Thus, this path integral representation takes into account quantum uncertainty of the quasiparticle position. In the limit of a large mass the spatial quasiparticle extension becomes much smaller than the average interparticle distance. This makes possible an analytical integration over the ‘beads’ positions by the method of steepest decent. As result the partition function is reduced to its classical limit involving point-like quasiparticles.

III. WIGNER DYNAMICS

We are going to use Wigner formulation of quantum mechanics for consideration of QGP kinetic properties. Let us review the underlying ideas of the Wigner dynamics for the simplest case, i.e. for nonrelativistic colorless system of particles [48]. The basis of our consideration is the Wigner representation of the von Neumann equation – a Wigner-Liouville equation (WLE). To derive the WLE for the simplest density matrix $\rho(r, r', t) = \Psi(r, t)\Psi^*(r', t)$ of the 3D N-particle system with Ψ being an eigenfunction of a Hamiltonian operator $\hat{H} = \sum_{i=1}^N \hat{p}_i^2/m + U$, we introduce center-of-mass and relative coordinates in a standard manner: $q \equiv (r + r')/2$ and $\xi \equiv r' - r$. Note that all these quantities are 3N-dimensional vectors. A Wigner distribution function (WF) is defined as

$$w(p, q, t) = \frac{1}{(2\pi)^{6N}} \int \rho\left(q + \frac{\xi}{2}, q - \frac{\xi}{2}, t\right) e^{ip\xi} d\xi. \quad (25)$$

Here and below products of vector quantities like $p\xi$ are understood as scalar products of 3N dimensional vectors. Using this definition, one can derive the WLE for $w(p, q, t)$ [48–50]. Applying time derivative to definition (25) and taking into account that

$$i\frac{\partial}{\partial t}\Psi(r, t) = \hat{H}\Psi(r, t), \quad i\frac{\partial}{\partial t}\Psi^*(r, t) = -\hat{H}\Psi^*(r, t) \quad (26)$$

we arrive at

$$\begin{aligned} \frac{\partial w(p, q, t)}{\partial t} &= \frac{1}{(2\pi)^{6N}} \int d\xi \exp(ip\xi) \frac{1}{i} [\hat{H}(r, t) - \hat{H}(r', t)] \rho(r, r', t) \\ &= \frac{1}{(2\pi)^{6N}} \int d\xi \exp(ip\xi) \left\{ -\frac{i}{m} \frac{\partial^2}{\partial q \partial \xi} + \frac{1}{i} \left[U\left(q - \frac{\xi}{2}\right) - U\left(q + \frac{\xi}{2}\right) \right] \right\} \rho\left(q + \frac{\xi}{2}, q - \frac{\xi}{2}, t\right). \end{aligned} \quad (27)$$

By means of integration by parts the first term in the braces can be transformed as follows

$$\frac{1}{(2\pi)^{6N}} \int d\xi \exp(ip\xi) \left(-\frac{i}{m}\right) \frac{\partial^2}{\partial q \partial \xi} \rho\left(q + \frac{\xi}{2}, q - \frac{\xi}{2}, t\right) = -\frac{p}{m} \frac{\partial w(p, q, t)}{\partial q} \quad (28)$$

while for the second one we obtain the following expression

$$\begin{aligned} & \frac{1}{(2\pi)^{6N}} \int d\xi \exp(ip\xi) \frac{1}{i} \left[U\left(q - \frac{\xi}{2}\right) - U\left(q + \frac{\xi}{2}\right) \right] \rho\left(q + \frac{\xi}{2}, q - \frac{\xi}{2}, t\right) \\ &= \frac{4}{(2\pi)^{6N}} \int ds w(p - s, q, t) \int dq' U(q - q') \sin(2sq') \end{aligned} \quad (29)$$

which results from substitution of ρ expressed in the form of inverse transformation to formula (25).

This way we arrive at the following form of the WLE

$$\frac{\partial w}{\partial t} + \frac{p}{m} \frac{\partial w}{\partial q} - \frac{\partial U(q)}{\partial q} \frac{\partial w}{\partial p} = \int ds w(p - s, q, t) \omega(s, q), \quad (30)$$

where

$$\omega(s, q) = -\frac{\partial U(q)}{\partial q} \frac{d\delta(s)}{ds} + \frac{4}{(2\pi)^{6N}} \int dq' U(q - q') \sin(2sq') \quad (31)$$

In the classical limit, $\hbar \rightarrow 0$, the r.h.s. of Eq. (30) disappears and Eq. (30) is reduced to the classical Liouville equation. This is the reason why we extracted the term $\partial U(q)/\partial q$ from the r.h.s. of Eq. (30).

A. Wigner Dynamics of Color Particles

Let us consider dynamics of QGP quasiparticles additionally characterized by color variables Q and derive the color WLE for a density matrix $\rho(r, r', Q, t)$ of the 3D N-particle system, where, as before, Q denotes color degrees of freedom of all quarks, antiquarks and gluons. Since color charges Q are treated classically, we consider only diagonal density matrix with respect to colors. Indeed, the Q variable already includes both canonical coordinate and momentum corresponding to classical color dynamics (see Appendix I). Therefore, the density matrix takes the form

$$\rho(r, r', Q, t) = \underline{\rho}(r, r', Q, t) \prod_i \delta(Q_i - Q_i(t)) \quad (32)$$

where the product runs over all particles in the system and $Q_i(t)$ is a solution of the classical equation of motion for color (4). Here $\underline{\rho}(r, r', Q, t) = \Psi(r, Q, t) \Psi^*(r', Q, t)$ is a quantum part of the density matrix with Ψ being an eigenfunction of the Hamiltonian operator described by Eq. (1) and Q are already fixed c-numbers.

Now the definition of the corresponding WF reads

$$w(p, q, Q, t) = \frac{1}{(2\pi)^{3N}} \int \rho\left(q - \frac{\xi}{2}, q + \frac{\xi}{2}, Q, t\right) e^{ip\xi} d\xi, \quad (33)$$

The quasiparticles are also characterized by spin and flavor, which we do not explicitly include in the list of quasiparticle degrees of freedom. Notice that color degrees of freedom are also in the Wigner representation, since Q includes both color canonical coordinated and momenta, see Appendix I. Now the WLE is defined by equation of the form [34, 51]:

$$\frac{\partial w}{\partial t} + v \frac{\partial w}{\partial q} + F \frac{\partial w}{\partial p} + \Upsilon \frac{\partial w}{\partial Q} = \int ds w(p - s, q, Q, t) \omega(s, q), \quad (34)$$

where $v = \{\mathbf{v}_i\}$ is 3N-dimensional vector of velocities of all quasiparticles, cf. Eq. (3), $F = -\partial U^C(q, Q)/\partial q$ is a set of the color-electric forces experienced by all quasiparticles, cf. Eq. (2), $\Upsilon = \{\Upsilon_i^a\}$ is an 8N-dimensional vector of driving forces in Wong's equation of motion for the color charge (4), and $\omega(s, q)$ is defined by Eq. (31).

The classical part of WLE (34), i.e. the l.h.s. of it, can be easily derived e.g. from the Wong's equations of motion (2)-(4) for the color-charged particles (see Ref. [51]). In particular, the term $\Upsilon \partial w / \partial Q$ naturally results from $(dQ(t)/dt) \partial w / \partial Q$ and Wong's equation (4). Since we confine ourselves to classical dynamics of color, we do not need any further (quantum) consideration for it. The quantum space dynamics [i.e. the r.h.s. of Eq. (34)] is derived completely in the same way as it was described above, see Eqs. (27)-(30), with minor complications due to relativistic kinematics.

B. Wigner representation of time correlation functions

In computations of transport properties, like viscosity, our starting point is the general Kubo expression for the canonical ensemble-averaged operator [52]

$$\check{C}_{BA}(t) = Z^{-1} \text{Tr} \left\{ e^{-\beta H} \hat{B} e^{i\hat{H}t} \hat{A} e^{-i\hat{H}t} \right\}, \quad (35)$$

where \hat{B} and \hat{A} are quantum operators of dynamic quantities under consideration and $Z(N, V, T) = \text{Tr} \left\{ e^{-\beta \hat{H}} \right\}$ is the canonical partition function. Frequently a symmetric time-correlation function is also used [53]:

$$C_{BA}(t) = Z^{-1} \text{Tr} \left\{ \hat{B} e^{i\hat{H}t_c} \hat{A} e^{-i\hat{H}t_c} \right\}, \quad (36)$$

where $t_c = t - i\beta/2$ is a complex-valued quantity including the inverse temperature $\beta = 1/T$. The Fourier transforms of $\check{C}_{BA}(t)$ and $C_{BA}(t)$ are related as [53]

$$C_{BA}(\omega) = \exp \left(-\frac{\beta\omega}{2} \right) \check{C}_{BA}(\omega). \quad (37)$$

As a consequence, transport coefficients described by zero-frequency ($\omega = 0$) Fourier components can be obtained from the symmetric time-correlation functions, which may offer certain computational advantages. This symmetric form is used below.

The Wigner representation of the time-correlation function in a $6N$ -dimensional space can be written as

$$C_{BA}(t) = (2\pi)^{-6N} \int d\overline{pq\mu Q} d\widetilde{pq\mu Q} B(\overline{pqQ}) A(\widetilde{pqQ}) W \left(\overline{pqQ}; \widetilde{pqQ}; t; \beta \right) \quad (38)$$

where we introduced a short-hand notation for phase space points in $(6N+8N)$ -dimensional space, \overline{pqQ} and \widetilde{pqQ} , with p, q and Q comprising the momenta, coordinates and color variables, respectively, of all particles of the system. Here $A(\widetilde{pqQ})$ denotes the Weyl's symbol [48] of the operator \hat{A} :

$$A(\widetilde{pqQ}) = \int d\tilde{\xi} \exp(-i\tilde{p}\tilde{\xi}) \left\langle \tilde{q} - \frac{\tilde{\xi}}{2}, \tilde{Q} \left| \hat{A} \right| \tilde{q} + \frac{\tilde{\xi}}{2}, \tilde{Q} \right\rangle, \quad (39)$$

and similarly for the operator \hat{B} , while $W \left(\overline{pqQ}; \widetilde{pqQ}; t; \beta \right)$ is the spectral density expressed as

$$W \left(\overline{pqQ}; \widetilde{pqQ}; t; \beta \right) = Z^{-1} \sum_{\tilde{\sigma}, \bar{\sigma}} \int \int d\tilde{\xi} d\bar{\xi} e^{i\tilde{p}\tilde{\xi}} e^{i\bar{p}\bar{\xi}} \left\langle \bar{q} + \frac{\bar{\xi}}{2}, \bar{Q} \left| e^{i\hat{H}t_c} \right| \tilde{q} - \frac{\tilde{\xi}}{2}, \tilde{Q} \right\rangle \left\langle \tilde{q} + \frac{\tilde{\xi}}{2}, \tilde{Q} \left| e^{-i\hat{H}t_c} \right| \bar{q} - \frac{\bar{\xi}}{2}, \bar{Q} \right\rangle. \quad (40)$$

In Eq. (38) we silently assumed that operators \hat{A} and \hat{B} do not depend on spin variables. Therefore, summation over spins $\tilde{\sigma}$ and $\bar{\sigma}$ can be safely moved to the definition of W . Here and below we do not explicitly write spin variables, if they are not essential. The time-correlation function $C_{BA}(t)$ is a linear functional of the spectral density W . Thus, the problem of its treatment is reduced to the consideration of the spectral density evolution.

As it follows from Eqs. (34) and Ref. [54], the following system of the WL integro-differential equations describe the time evolution of the color spectral density W :

$$\frac{\partial W}{\partial t} + \tilde{v} \frac{\partial W}{\partial \tilde{q}} + F \frac{\partial W}{\partial \tilde{p}} + \Upsilon \frac{\partial W}{\partial \tilde{Q}} = \int ds W \left(\tilde{p} - s, \tilde{qQ}; \widetilde{pqQ}; t; \beta \right) \omega(s, \tilde{q}), \quad (41)$$

$$-\frac{\partial W}{\partial t} + \bar{v} \frac{\partial W}{\partial \bar{q}} + F \frac{\partial W}{\partial \bar{p}} + \Upsilon \frac{\partial W}{\partial \bar{Q}} = \int ds W \left(\overline{pqQ}; \tilde{p} - s, \bar{qQ}; t; \beta \right) \omega(s, \bar{q}), \quad (42)$$

where as before $\omega(s, q)$ is defined by Eq. (31). This equations are derived precisely in the same way as those of Eqs. (34) and (27)-(30), only the Hamiltonian H appears here as a result of time derivation of exponent functions, $e^{i\hat{H}t}$ and $e^{-i\hat{H}t}$, rather than from application of equations of motion (26) in Eq. (27). Notice that while Eq. (41) describes evolution in the positive time direction, Eq. (42) specifies propagation in the reverse time direction. This happens because of the presence the direct time $e^{-i\hat{H}t}$ and reverse time $e^{i\hat{H}t}$ evolution operators in definition of the time-correlation function (36).

Now using Eqs. (41) and (42) we can obtain an integral equation [48–50, 54] for $W(\overline{pqQ}; \widetilde{pqQ}; t; \beta)$

$$\begin{aligned}
& W(\overline{pqQ}; \widetilde{pqQ}; t; \beta) \\
&= \left\{ \int d\overline{p_0 q_0 \mu Q_0} d\overline{p_0 q_0 \mu Q_0} G(\overline{pqQ}, \widetilde{pqQ}, t; \overline{p_0 q_0 Q_0}, \widetilde{p_0 q_0 Q_0}, 0) W(\overline{p_0 q_0 Q_0}; \widetilde{p_0 q_0 Q_0}; t=0, \beta) \right. \\
&+ \frac{1}{2} \left(\int_0^t dt' \int ds \int d\overline{p' q' \mu Q'} d\overline{p' q' \mu Q'} G(\overline{pqQ}, \widetilde{pqQ}, t; \overline{p' q' Q'}, \widetilde{p' q' Q'}, t') \right. \\
&\times \left. \left[W(\overline{p'} - s, \overline{q' Q'}; \widetilde{p' q' Q'}; t'; \beta) \omega(s, \overline{q'}) - W(\overline{p' q' Q'}; \widetilde{p'} - s, \widetilde{q' Q'}; t'; \beta) \omega(s, \widetilde{q'}) \right] \right\}, \tag{43}
\end{aligned}$$

with Green function

$$\begin{aligned}
& G(\overline{pqQ}, \widetilde{pqQ}, t; \overline{p' q' Q'}, \widetilde{p' q' Q'}, t') \\
&= \delta(\overline{p} - \overline{p}(t; \overline{p' q' Q'}, t')) \delta(\overline{q} - \overline{q}(t; \overline{p' q' Q'}, t')) \delta(\overline{Q} - \overline{Q}(t; \overline{p' q' Q'}, t')) \\
&\times \delta(\widetilde{p} - \widetilde{p}(t; \widetilde{p' q' Q'}, t')) \delta(\widetilde{q} - \widetilde{q}(t; \widetilde{p' q' Q'}, t')) \delta(\widetilde{Q} - \widetilde{Q}(t; \widetilde{p' q' Q'}, t')). \tag{44}
\end{aligned}$$

describing propagation of the spectral density along classical trajectories in positive time direction

$$\begin{aligned}
\frac{d\overline{p}(t; \overline{p' q' Q'}, t')}{dt} &= \frac{1}{2} F(\overline{qQ}_t), \\
\frac{d\overline{q}(t; \overline{p' q' Q'}, t')}{dt} &= \frac{1}{2} \overline{v}[\overline{p}(t; \overline{p' q' Q'}, t')], \\
\frac{d\overline{Q}(t; \overline{p' q' Q'}, t')}{dt} &= \frac{1}{2} \Upsilon(\overline{qQ}_t), \tag{45}
\end{aligned}$$

and in the reverse time direction

$$\begin{aligned}
\frac{d\widetilde{p}(t; \widetilde{p' q' Q'}, t')}{dt} &= -\frac{1}{2} F(\widetilde{qQ}_t), \\
\frac{d\widetilde{q}(t; \widetilde{p' q' Q'}, t')}{dt} &= -\frac{1}{2} \widetilde{v}[\widetilde{p}(t; \widetilde{p' q' Q'}, t')], \\
\frac{d\widetilde{Q}(t; \widetilde{p' q' Q'}, t')}{dt} &= -\frac{1}{2} \Upsilon(\widetilde{qQ}_t), \tag{46}
\end{aligned}$$

where $(\widetilde{qQ}_t) = [\widetilde{q}(t; \widetilde{p' q' Q'}, t'), \widetilde{Q}(t; \widetilde{p' q' Q'}, t')]$ and similarly for bared quantities. These equations of motion are supplemented by initial conditions at time $t = 0$

$$\overline{p}(t; \overline{p_0 q_0 Q_0}, 0) = \overline{p_0}, \quad \overline{q}(t; \overline{p_0 q_0 Q_0}, 0) = \overline{q_0}, \quad \overline{Q}(t; \overline{p_0 q_0 Q_0}, 0) = \overline{Q_0}, \tag{47}$$

$$\widetilde{p}(t; \widetilde{p_0 q_0 Q_0}, 0) = \widetilde{p_0}, \quad \widetilde{q}(t; \widetilde{p_0 q_0 Q_0}, 0) = \widetilde{q_0}, \quad \widetilde{Q}(t; \widetilde{p_0 q_0 Q_0}, 0) = \widetilde{Q_0}. \tag{48}$$

and by initial conditions at time $t = t'$

$$\overline{p}(t'; \overline{p' q' Q'}, t') = \overline{p'}, \quad \overline{q}(t'; \overline{p' q' Q'}, t') = \overline{q'}, \quad \overline{Q}(t'; \overline{p' q' Q'}, t') = \overline{Q'}, \tag{49}$$

$$\widetilde{p}(t'; \widetilde{p' q' Q'}, t') = \widetilde{p'}, \quad \widetilde{q}(t'; \widetilde{p' q' Q'}, t') = \widetilde{q'}, \quad \widetilde{Q}(t'; \widetilde{p' q' Q'}, t') = \widetilde{Q'}. \tag{50}$$

In fact, Eqs. (45) are Wong's equations of motion but written for half-time ($t/2$). Similarly, Eqs. (46) are half-time Wong's equations of motion reversed in time. This happens because the time correlation is taken between instants in the past and the future with the initial conditions fixed in between these instants, i.e. at $t = 0$ the spectral density $W(\overline{pqQ}_0; \widetilde{pqQ}_0; t=0, \beta) = W_0(\overline{pqQ}_0; \widetilde{pqQ}_0; \beta)$ is fixed as it is described in the next subsection.

Solution of the integral equation (43) can be obtained in a form of iterative series. In this work, we take into account only the first term of this iteration series:

$$\begin{aligned}
& W(\overline{pqQ}; \widetilde{pqQ}; t; \beta) \\
&\simeq \int d\overline{p_0 q_0 \mu Q_0} d\overline{p_0 q_0 \mu Q_0} G(\overline{pqQ}, \widetilde{pqQ}, t; \overline{p_0 q_0 Q_0}, \widetilde{p_0 q_0 Q_0}, 0) W_0(\overline{pqQ}_0; \widetilde{pqQ}_0; \beta). \tag{51}
\end{aligned}$$

Notice that if the initial $W_0(\overline{pqQ}_0; \widetilde{pqQ}_0; \beta)$ is chosen appropriately [48], i.e. such that it contains all powers of the Planck's constant, then the first term of the iterative series, i.e. Eq. (51), describes propagation of a *quantum* initial spectral density along classical trajectories. Other (higher) terms describe propagation of the initial spectral density along the analogous trajectories but perturbed by momentum jumps resulted from the convolution structure of integral term in Eq. (43). From physical point of view these jumps relate to quantum $([p, q])$ uncertainty (for details see discussion in Ref. [49]). As it was found in Refs. [49, 50, 55], the main contribution to WF comes from the trajectories without jumps, i.e. from the first term of the iterative series, if the motion takes place in a classically accessible region, which is the case here. Nevertheless calculations with higher-order iterative terms are in progress and will be reported elsewhere.

Moreover, we replace the color Coulomb potential U^C by the Kelbg one Φ in quantities F and Υ defining propagation of the spectral density along classical trajectories, see Eqs. (45) and (46). By this replacement we are able to take into account certain higher-order quantum terms of the iteration series presenting the solution of integral equation (43). From the practical point of view, it allows us to avoid problems due to singular character of the Coulomb potential. As known, the classical limit for multi-component Coulomb system does not exist since the stability of Coulomb systems is only provided by quantum effects. That is the physical reason for arising the color Kelbg potential in the partition function and Eq. (61) for initial condition (see below).

C. Initial conditions

The initial function W_0 is expressed in terms of matrix elements of density matrix considered in section II B. Accordingly to Eq. (40) for $t = 0$ and definition of the density matrix ρ , cf. Eq. (7), we have

$$W_0(\overline{pqQ}; \widetilde{pqQ}; \beta) = Z^{-1} \sum_{\bar{\sigma}, \tilde{\sigma}} \int d\bar{\xi} d\tilde{\xi} e^{i\bar{p}\bar{\xi}} e^{i\tilde{p}\tilde{\xi}} \left\langle \bar{q} + \frac{\bar{\xi}}{2}, \bar{Q} \left| \rho\left(\frac{\beta}{2}\right) \right| \tilde{q} - \frac{\tilde{\xi}}{2}, \tilde{Q} \right\rangle \left\langle \tilde{q} + \frac{\tilde{\xi}}{2}, \tilde{Q} \left| \rho\left(\frac{\beta}{2}\right) \right| \bar{q} - \frac{\bar{\xi}}{2}, \bar{Q} \right\rangle \times \delta(\bar{Q} - \tilde{Q}) \delta_{\bar{\sigma}, \tilde{\sigma}}. \quad (52)$$

Thus, the problem is reduced to calculation of matrix elements of density matrix ρ , which is similar to that we did in sect. II devoted to thermodynamics. Only now we need nondiagonal matrix elements rather than diagonal ones, as in sect. II.

As before, cf. Eq. (12), let us subdivide $\rho(\beta/2)$ into beards using the operator identity

$$e^{-\beta \hat{H}/2} = e^{-\Delta\beta' \hat{H}} \cdot e^{-\Delta\beta' \hat{H}} \dots e^{-\Delta\beta' \hat{H}},$$

where the r.h.s. contains $n + 1$ identical factors with $\Delta\beta' = \beta/[2(n + 1)]$, so

$$\left\langle \bar{q} + \frac{\bar{\xi}}{2}, Q \left| \rho\left(\frac{\beta}{2}\right) \right| \tilde{q} - \frac{\tilde{\xi}}{2}, Q \right\rangle \simeq \int d\bar{r}^{(1)} \dots d\bar{r}^{(n)} \bar{\rho}^{(1)} \bar{\rho}^{(2)} \bar{\rho}^{(3)} \dots \bar{\rho}^{(n)} \bar{\rho}^{(n+1)} \quad (53)$$

where $\bar{\rho}^{(l)}$ ($l = 1, \dots, n + 1$) are defined by Eqs. (13) and (19) with

$$\bar{r}^{(0)} = \bar{q} + \frac{\bar{\xi}}{2}, \quad \bar{r}^{(n+1)} = \tilde{q} - \frac{\tilde{\xi}}{2}$$

and $\Delta\beta$ replaced by $\Delta\beta'$. The bar sign above $\bar{\rho}^{(l)}$ means that these quantities depend on “bared” variables $\bar{r}^{(l)}$. Similarly, after antisymmetrization⁴ we obtain

$$\left\langle \tilde{q} + \frac{\tilde{\xi}}{2}, Q \left| \rho\left(\frac{\beta}{2}\right) \right| \bar{q} - \frac{\bar{\xi}}{2}, Q \right\rangle \simeq \int d\bar{r}^{(1)} \dots d\bar{r}^{(n+1)} \bar{\rho}^{(1)} \bar{\rho}^{(2)} \bar{\rho}^{(3)} \dots \bar{\rho}^{(n)} \times \left[\sum_{P_q} \sum_{P_{\bar{q}}} \sum_{P_g} (-1)^{\kappa_{P_q} + \kappa_{P_{\bar{q}}}} \hat{P}_q \hat{P}_{\bar{q}} \hat{P}_g \bar{\rho}^{(n+1)} \right], \quad (54)$$

⁴ Symmetrization-antisymmetrization is performed only in one matrix element in Eq. (52), since it is enough to make the full expression under the trace (38) to be symmetrized-antisymmetrized.

with

$$\tilde{r}^{(0)} = \tilde{q} + \frac{\tilde{\xi}}{2}, \quad \tilde{r}^{(n+1)} = \tilde{q} - \frac{\tilde{\xi}}{2},$$

where the “tilde” functions $\tilde{\rho}^{(l)}$ depend on “tilde” variables.

In Sect. II we used the approximate expression for elements of the density matrix defined by Eq. (19). Here it is not practical. An alternative approximation can be derived by means of the following expression for the density matrix operator

$$\exp(-\Delta\beta'\hat{H}) \approx \exp\left(-\frac{\Delta\beta'}{2}\hat{K}\right) \exp(-\Delta\beta'\hat{U}^C) \exp\left(-\frac{\Delta\beta'}{2}\hat{K}\right) \left\{ \text{terms with } [\hat{K}, \hat{U}^C] \right\}, \quad (55)$$

compare with Eq. (14). Notice that in fact $\exp\{-(\Delta\beta'/2)\hat{K}\}$ is a density matrix of the non-interacting system. This symmetrized form results in the following approximation to the matrix elements

$$\begin{aligned} \rho^{(l)} &= \rho\left(r^{(l-1)}, r^{(l)}, Q; \{N\}; \Delta\beta'/2\right) \\ &= \int dq^{(l)} \rho_0\left(r^{(l-1)}, q^{(l)}, Q; \{N\}; \Delta\beta'/2\right) \rho_0\left(q^{(l)}, r^{(l)}, Q; \{N\}; \Delta\beta'/2\right) \\ &\times \exp\left[-\Delta\beta'\frac{1}{2} \sum_{i,j(i \neq j)}^N \Phi_{ij}\left(\mathbf{q}_i^{(l)} - \mathbf{q}_j^{(l)}, \mathbf{q}_i^{(l)} - \mathbf{q}_j^{(l)}, Q_i, Q_j\right)\right], \end{aligned} \quad (56)$$

where Φ_{ij} is the diagonal part of the color Kelbg potential, see Eq. (18). This approximate form has the same accuracy as that in Eq. (13). At the same time it allows us to explicitly perform Fourier transforms in Eq. (52), since the ξ dependence now occurs only in the ρ_0 factors.

Thus, based on the above approximation for $\rho^{(l)}$ we are able to explicitly evaluate integrals over $\tilde{\xi}$ and $\tilde{\xi}$

$$\int d\tilde{\xi} e^{i\tilde{p}\tilde{\xi}} \tilde{\rho}^{(n+1)} \tilde{\rho}^{(1)} = \tilde{\rho}^{(n+1)} \tilde{\rho}^{(1)} \varphi\left(\tilde{p}; \tilde{r}^{(n)}, \tilde{r}^{(1)}\right) \quad (57)$$

$$\int d\tilde{\xi} e^{i\tilde{p}\tilde{\xi}} \tilde{\rho}^{(n+1)} \tilde{\rho}^{(1)} = \tilde{\rho}^{(n+1)} \tilde{\rho}^{(1)} \varphi\left(\tilde{p}; \tilde{r}^{(n)}, \tilde{r}^{(1)}\right) \quad (58)$$

where on the r.h.s. of these equations and below the marginal coordinates take already the following values

$$\bar{r}^{(0)} = \bar{q}, \quad \bar{r}^{(n+1)} = \tilde{q},$$

$$\tilde{r}^{(0)} = \tilde{q}, \quad \tilde{r}^{(n+1)} = \bar{q},$$

and the complex-valued function φ is defined as

$$\varphi(p; r', r'') = \prod_{i=1}^N (2\lambda_i'^2)^{3/2} \exp\left[-\frac{1}{2\pi} \left(\mathbf{p}_i \lambda_i' + i\pi \frac{\mathbf{r}_i' - \mathbf{r}_i''}{\lambda_i'}\right)^2\right], \quad (59)$$

with $\lambda_i' = \sqrt{\pi\Delta\beta'/m_i}$ being the i -particle thermal wave lengths related to temperature $2/\Delta\beta'$.

Substituting these expressions into Eq. (52), we arrive at [46, 54, 55]

$$\begin{aligned} W_0(\overline{pqQ}; \widetilde{pqQ}; \beta) &= \frac{1}{Z} \int d\bar{r}^{(1)} \dots d\bar{r}^{(n)} d\tilde{q}^{(1)} \dots d\tilde{q}^{(n+1)} d\tilde{r}^{(1)} \dots d\tilde{r}^{(n)} d\tilde{q}^{(1)} \dots d\tilde{q}^{(n+1)} \\ &\times \Psi\left(\overline{pqQ}; \widetilde{pqQ}; \bar{r}^{(1)}, \dots, \bar{r}^{(n)}; \bar{q}^{(1)}, \dots, \bar{q}^{(n+1)}; \tilde{r}^{(1)}, \dots, \tilde{r}^{(n)}; \tilde{q}^{(1)}, \dots, \tilde{q}^{(n+1)}; \beta\right) \end{aligned} \quad (60)$$

with

$$\begin{aligned} &\Psi\left(\overline{pqQ}; \widetilde{pqQ}; \bar{r}^{(1)}, \dots, \bar{r}^{(n)}; \bar{q}^{(1)}, \dots, \bar{q}^{(n+1)}; \tilde{r}^{(1)}, \dots, \tilde{r}^{(n)}; \tilde{q}^{(1)}, \dots, \tilde{q}^{(n+1)}; \beta\right) \\ &= \exp\left\{-\beta\left(\bar{U} + \tilde{U}\right)\right\} \delta(\bar{Q} - \tilde{Q}) \end{aligned}$$

$$\begin{aligned}
& \times \sum_{\bar{\sigma}, \bar{\sigma}'} \left[\prod_{l=1}^{n+1} \rho_0 \left(\bar{r}^{(l-1)}, \bar{q}^{(l)}, \bar{Q}; \{N\}; \Delta\beta'/2 \right) \rho_0 \left(\bar{q}^{(l)}, \bar{r}^{(l)}, \bar{Q}; \{N\}; \Delta\beta'/2 \right) \right] \varphi \left(\bar{p}; \bar{r}^{(n)}, \bar{r}^{(1)} \right) \\
& \times \left[\prod_{l=1}^n \rho_0 \left(\tilde{r}^{(l-1)}, \tilde{q}^{(l)}, \tilde{Q}; \{N\}; \Delta\beta'/2 \right) \rho_0 \left(\tilde{q}^{(l)}, \tilde{r}^{(l)}, \tilde{Q}; \{N\}; \Delta\beta'/2 \right) \right] \varphi \left(\bar{p}; \tilde{r}^{(n)}, \tilde{r}^{(1)} \right) \\
& \times \left[\sum_{P_q} \sum_{P_{\bar{q}}} \sum_{P_g} (-1)^{\kappa_{P_q} + \kappa_{P_{\bar{q}}}} \hat{P}_q \hat{P}_{\bar{q}} \hat{P}_g \rho_0 \left(\tilde{q}^{(n+1)}, \tilde{r}^{(n+1)}, \tilde{Q}; \{N\}; \Delta\beta'/2 \right) \delta_{\bar{\sigma}, \bar{\sigma}'} \mathcal{S}(\bar{\sigma}, \hat{P}_q \hat{P}_{\bar{q}} \hat{P}_g \bar{\sigma}') \Big|_{\bar{\sigma}' = \bar{\sigma}} \right] \quad (61)
\end{aligned}$$

where

$$\bar{U} = \frac{1}{2} \frac{1}{n+1} \sum_{l=1}^{n+1} \sum_{i,j(i \neq j)}^N \Phi_{ij} \left(\bar{\mathbf{q}}_i^{(l)} - \bar{\mathbf{q}}_j^{(l)}, \bar{\mathbf{q}}_i^{(l)} - \bar{\mathbf{q}}_j^{(l)}, \bar{Q}_i, \bar{Q}_j \right). \quad (62)$$

$$\tilde{U} = \frac{1}{2} \frac{1}{n+1} \sum_{l=1}^{n+1} \sum_{i,j(i \neq j)}^N \Phi_{ij} \left(\tilde{\mathbf{q}}_i^{(l)} - \tilde{\mathbf{q}}_j^{(l)}, \tilde{\mathbf{q}}_i^{(l)} - \tilde{\mathbf{q}}_j^{(l)}, \tilde{Q}_i, \tilde{Q}_j \right). \quad (63)$$

Applying the notation of Sect. II (cf. Eqs. (22) and Eq. (23)), we finally arrive at

$$\begin{aligned}
& \Psi \left(\overline{pqQ}; \widetilde{pqQ}; r^{(1)}, \dots, \bar{r}^{(n)}; \bar{q}^{(1)}, \dots, \bar{q}^{(n+1)}; \tilde{r}^{(1)}, \dots, \tilde{r}^{(n)}; \tilde{q}^{(1)}, \dots, \tilde{q}^{(n+1)}; \beta \right) \\
& = \exp \left\{ -\beta \left(\bar{U} + \tilde{U} \right) \right\} \delta(\bar{Q} - \tilde{Q}) \\
& \times \sum_{\bar{\sigma}, \bar{\sigma}'} \left[\prod_{l=1}^{n+1} \left(\prod_{i=1}^N \phi_{ii} \left(\bar{r}^{(l-1)}, \bar{q}^{(l)}, \Delta\beta'/2 \right) \right) \left(\prod_{i=1}^N \phi_{ii} \left(\bar{q}^{(l)}, \bar{r}^{(l)}, \Delta\beta'/2 \right) \right) \right] \varphi \left(\bar{p}; \bar{r}^{(n)}, \bar{r}^{(1)} \right) \\
& \times \left[\prod_{l=1}^n \left(\prod_{i=1}^N \phi_{ii} \left(\tilde{r}^{(l-1)}, \tilde{q}^{(l)}, \Delta\beta'/2 \right) \right) \left(\prod_{i=1}^N \phi_{ii} \left(\tilde{q}^{(l)}, \tilde{r}^{(l)}, \Delta\beta'/2 \right) \right) \right] \\
& \times \frac{\det \|\phi(\bar{q}^{(n+1)}, \tilde{r}^{(n+1)}, \Delta\beta'/2)\|_{N_q}}{\Lambda_q^{6(n+1)N_q}(\Delta\beta'/2)} \frac{\det \|\phi(\tilde{q}^{(n+1)}, \bar{r}^{(n+1)}, \Delta\beta'/2)\|_{\bar{N}_q}}{\Lambda_{\bar{q}}^{6(n+1)\bar{N}_q}(\Delta\beta'/2)} \\
& \times \frac{\text{per} \|\phi(\tilde{q}^{(n+1)}, \bar{r}^{(n+1)}, \Delta\beta'/2)\|_{N_g}}{\Lambda_g^{6(n+1)N_g}(\Delta\beta'/2)} \varphi \left(\bar{p}; \tilde{r}^{(n)}, \tilde{r}^{(1)} \right) \delta_{\bar{\sigma}, \bar{\sigma}'} \quad (64)
\end{aligned}$$

with

$$\bar{r}^{(0)} = \bar{q}, \quad \bar{r}^{(n+1)} = \tilde{q}, \quad \tilde{r}^{(0)} = \tilde{q}, \quad \tilde{r}^{(n+1)} = \bar{q}.$$

In the limit $n \rightarrow \infty$ this expression exactly gives the product of matrix elements in Eq. (52) in the form of path integrals multiplied by a limiting expression of the φ functions. According to the Lebesgue-Dirac delta theorem (see Appendix II), the $\bar{\rho}^{(0)} \tilde{\rho}^{(0)} \varphi$ products in integral (60) in the limit $n \rightarrow \infty$ are equivalent to the following real valued expressions

$$\bar{\rho}^{(n+1)} \tilde{\rho}^{(1)} \varphi \left(\bar{p}; \bar{r}^{(n)}, \bar{r}^{(1)} \right) \xrightarrow{n \rightarrow \infty} \bar{\rho}^{(n+1)} \tilde{\rho}^{(1)} \prod_{i=1}^N \left((2\lambda_i^2)^{3/2} \exp \left[-\frac{\tilde{\mathbf{p}}_i^2 \lambda_i^2}{2\pi} \right] \right) \left(\frac{\lambda_i}{\pi} \right)^3 \delta \left(\bar{\mathbf{r}}_i^{(n)} - \tilde{\mathbf{r}}_i^{(1)} \right) \quad (65)$$

$$\tilde{\rho}^{(n+1)} \bar{\rho}^{(1)} \varphi \left(\bar{p}; \tilde{r}^{(n)}, \tilde{r}^{(1)} \right) \xrightarrow{n \rightarrow \infty} \tilde{\rho}^{(n+1)} \bar{\rho}^{(1)} \prod_{i=1}^N \left((2\lambda_i^2)^{3/2} \exp \left[-\frac{\bar{\mathbf{p}}_i^2 \lambda_i^2}{2\pi} \right] \right) \left(\frac{\lambda_i}{\pi} \right)^3 \delta \left(\tilde{\mathbf{r}}_i^{(n)} - \bar{\mathbf{r}}_i^{(1)} \right) \quad (66)$$

where $\lambda_i = \sqrt{\pi\beta/2m_i}$. Analytical integration over the delta function simplifies the final path integral used for further computation of the real-valued W_0 by means of the Monte-Carlo method.

IV. SIMULATIONS OF QGP

The developed approach is applied to the QGP at zero baryon density ($\mu_q = 0$). Then assumption on equal quark masses (see point **IV** in subsect. II A) immediately implies equal fractions of quarks and antiquarks of different flavors: $N_u = N_d = N_s = N_q/3 = \bar{N}_u = \bar{N}_d = \bar{N}_s = \bar{N}_q/3$. Ideally the parameters of the model should be deduced from the QCD lattice data. However, presently this task is still quite ambiguous. Therefore, in the present simulations we take only a possible set of parameters. We use so called “one-loop analytic coupling” [56, 57]

$$\alpha_s(Q^2) = \frac{4\pi}{11 - (2/3)N_f} \left[\frac{1}{\ln(Q^2/\Lambda_{QCD}^2)} + \frac{\Lambda_{QCD}^2}{\Lambda_{QCD}^2 - Q^2} \right] \quad (67)$$

where Q is the momentum transfer, $\Lambda_{QCD} = 206$ MeV is the QCD scale and $N_f = 3$ is the number of flavors. The analytically generated non-perturbative contribution $\Lambda_{QCD}^2/(\Lambda_{QCD}^2 - Q^2)$ subtracts the unphysical Landau pole in a minimal way, yielding a ghost-free behavior which avoids any adjustable parameter. This coupling agrees with a great body of experimental data [56, 57]. As it is usually done in thermal field models, we substitute Q by $2\pi T$ to use this coupling in our simulations. The resulting $\alpha_s(T)$ is displayed in the left panel of Fig. 1 and compared with QCD-lattice coupling deduced from a short-distance behavior of the singlet free energy [58] and from spectral density of heavy-quark correlator [59]. As seen, the running coupling deduced from experimental data is close to those obtained in the lattice QCD. Notice that determination of $\alpha_s(T)$ in lattice QCD simulations is quite indirect. Therefore, different indirect methods naturally give somewhat different results.




FIG. 1: Input data for the PIMC simulations.

Left panel: Running coupling constant versus temperature fitted to experimental data [56, 57] (solid line). Points present the coupling deduced from lattice QCD simulations in Refs. [58] [lattice (Bielefeld 2012)] and [59] [lattice (Banerjee et al. 2012)]. Right panel: Mass-to-temperature ratio for quark and gluon quasiparticles versus temperature. Points are values used in simulations. The solid lines are smooth interpolations between points.

The quasiparticle masses were chosen to reproduce the pressure obtained in lattice QCD calculations [4, 5]. The T -dependence of these masses is presented in Fig. 1(right panel). When choosing masses we kept in mind constraints resulting from lattice QCD data [35, 37, 60] and QCD-motivated quasiparticle models [61, 62]. While gluon masses used in this paper well comply with those deduced from both lattice QCD data [35, 60] and quasiparticle models, this is not the case for quark masses. Our quark masses agrees with values required for quasiparticle fits [61, 62] of the lattice thermodynamic properties of the QGP: $m_q/T \simeq 1.5 \div 2.5$. At the same time they are appreciably lower than those in old lattice data [35]: $m_q/T \simeq 4$, and higher than $m_q/T \simeq 0.8$ reported in newer lattice calculations [37].

A. Equilibrium Properties

Figure 2 (left panel) demonstrates the quality of reproduction of the equation of state (EOS), i.e. the pressure versus temperature, achieved with the above discussed input data. The reference EoS (filled points in the left panel of Fig. 2) is taken from QCD lattice simulations of the QGP [4]. The quality of the reproduction obviously depends on the degree of accurate tuning of the input data, i.e. the quasiparticle masses. However, the PIMC scheme itself also produces certain errors. If there are metastable states of the system, convergence of calculations becomes poor

because of jumps between stable and metastable states. This is a typical situation when the system approaches to a point (or a range) of a phase transition. Precisely this happens at the lower end of considered temperature range. The shaded area in the left panel of Fig. 2 indicates these uncertainties of the PIMC calculations. Figure 2 also presents the entropy S/T^3 and trace anomaly $(\varepsilon - 3P)/T^4$ of the QGP. These quantities are calculated accordingly to Eqs. (8)-(11). In order to avoid the numeric noise, the derivative of a smooth interpolation between the PIMC points (solid line in the left panel of Fig. 2) was taken. Though agreement with the lattice data looks worse for the entropy and trace anomaly, in fact it is the same as that for pressure. Differentiation operations in Eqs. (9) and (11) make differences between PIMC results and lattice data more pronounced.




FIG. 2: Pressure (left panel), entropy (middle panel) and trace anomaly (right panel) scaled with corresponding powers of temperature versus temperature from PIMC simulations (open squares). These are compared with lattice data of Refs. [4, 5] (filled circles). The solid line in the left panel is a smooth interpolation between the PIMC points. Entropy and trace anomaly are calculated based on this smooth interpolation. The shaded area in the left panel indicates uncertainties of the PIMC calculations.

Having calibrated the model by reproducing the EoS, we can proceed to predictions. First, let us consider internal properties of the system. To characterize physical conditions and interplay of interaction and degeneracy in Fig. 3 a degeneracy parameter χ_u for 'up' quarks and a plasma coupling parameter Γ are presented:

$$\chi_u = n_u \lambda_u^3, \quad \Gamma = \frac{\bar{q}_2 g^2}{4\pi r_s T}, \quad (68)$$

where the thermal wave length λ_u was defined in the previous sections (see text after Eq. (22)), n_u is density of u quarks, $r_s^3 = 3/(4\pi n)$ is Wigner-Seitz radius, n is the density of all quasiparticles (quarks, antiquarks and gluons), and \bar{q}_2 is the quadratic Casimir value averaged over quarks, antiquarks and gluons, $\bar{q}_2 = N_c^2 - 1$ is a good estimate for this quantity. The plasma coupling parameter is a measure of ratio of the average potential to the average kinetic energy, and the degeneracy parameter χ_u indicates wheather a system is classical ($\chi_u \ll 1$) or quantum ($\chi_u \gtrsim 1$). It turns out that Γ and χ_u are of order unity which indicates that the QGP is a strongly coupled *quantum liquid* ($\Gamma \sim 1$) rather than a gas.

To clarify interplay of interaction and degeneracy let us consider spatial arrangement of the quasiparticles in the QGP by studying a pair distribution function (PDF) $g_{ab}(R)$ defined as

$$g_{ab}(|\mathbf{R}_1 - \mathbf{R}_2|) = \left(\frac{V}{N}\right)^2 \sum_{\sigma} \sum_{i,j,i \neq j} \delta_{a_i,a} \delta_{a_j,b} \frac{1}{Z} \int dr d\mu Q \delta(\mathbf{R}_1 - \mathbf{r}_i) \delta(\mathbf{R}_2 - \mathbf{r}_j) \rho(r, Q, \sigma; \{N\}; \beta), \quad (69)$$

where a_i and a_j are types of the particles ($= q, \bar{q}$ or g). The PDF gives a probability density to find a pair of particles of types a and b at a certain distance $R = |\mathbf{R}_1 - \mathbf{R}_2|$ from each other. The PDF depends only on the difference of coordinates because of the translational invariance of the system. In a non-interacting classical system, $g_{ab}(R) \equiv 1$, whereas interactions and quantum statistics result in a redistribution of the particles. At temperatures $T = 525$ MeV and $T = 193$ MeV the PDF averaged over the quasiparticle spin, colors and flavors are shown in Fig. 4.

At distances $R \gtrsim 0.2$ or 0.3 fm, depending on the temperature, all PDF's are practically equal to unity (Fig. 4) like in ideal gas due to the screening of the color Coulomb interaction. A drastic difference between qq and gg PDF's (the $\bar{q}\bar{q}$ PDF is identical to the qq one) occurs at short distances. Here the gluon-gluon and gluon-quark PDF's increase monotonically when the distance goes to zero while the qq and $\bar{q}\bar{q}$ ones remain uncorrelated. One of the physical reasons of the PDF difference is spatial quantum uncertainty and different properties of Bose and Fermi statistics of gluon and (anti)quark quasiparticles. Uncertainty in particle localization is defined by the ratio T/m . Localization




FIG. 3: Quark degeneracy parameter χ_u and the plasma coupling parameter Γ [see Eq. (68)] versus temperature. Lines are smooth interpolations between points.




FIG. 4: Pair correlation functions of identical (left panels) and different (right panels) quasiparticles at temperatures $T = 525$ MeV (upper panels) and $T = 193$ MeV (lower panels).

is better for heavier gluon quasiparticles. Fermi statistics results in effective quark-quark and antiquark-antiquark repulsion, while Bose one results in effective gluon-gluon attraction. Oscillations of the PDF at very small distances $R \lesssim 0.1$ fm are related to Monte-Carlo statistical error, as probability of quasiparticles being at short distances quickly decreases.

However, the qq and $\bar{q}\bar{q}$ pair correlation functions reveal practical absence of fermi repulsion. This happens because another physical reason comes into play. Strong interaction between quasiparticles reduces the influence of the degeneracy in the region of $\chi_u \sim 1$. This interaction is dominated by attraction at short distances. Indeed, the QGP lowers its total energy by minimizing the color Coulomb interaction energy via a spontaneous “anti-ferromagnetic”-




FIG. 5: Gluon-gluon and quark-antiquark pair correlation functions multiplied by R^2 at $T = 193$ MeV.

like ordering of color vectors, i.e. the color vectors of nearest neighbor quasiparticles become anti-parallel. Similar absence of fermi repulsion was observed in hydrogen plasma at $\chi \sim 1$ [47]. This short-distance attraction is stronger for gluon-gluon and gluon-(anti)quark pairs than for (anti)quark-(anti)quark ones because of the corresponding difference in values of quadratic Casimir invariants \check{q}_2 (see Appendix I), which determine the maximal values of the effective color charge products $|Q_i \cdot Q_j|$ in color Kelbg (Coulomb) potentials: For gluon-gluon pairs $|Q_g \cdot Q_g|_{max} = 24$, for gluon-(anti)quark pairs $|Q_g \cdot Q_q|_{max} = |Q_g \cdot Q_{\bar{q}}|_{max} \approx 10$, and for (anti)quark-(anti)quark pairs $|Q_q \cdot Q_q|_{max} = |Q_{\bar{q}} \cdot Q_{\bar{q}}|_{max} = |Q_q \cdot Q_{\bar{q}}|_{max} = 4$. Stronger gg attraction additionally enhances correlation of the gluon-gluon pairs at short distances. At the same time the short-distance attraction is the only reason of the gluon-(anti)quark short-distance correlation.

The short-distance correlation implies formation of the gluon-gluon and gluon-(anti)quark clusters, which are uniformly distributed in space. In case of the gluon-gluon clusters we can even speak about gg bound states, i.e. glueballs, as it is seen from Fig. 5. The product $R^2 g_{ab}(R)$ is proportional (up to constant factor) to a probability to find a pair of quasiparticles at a distance R from each other. As known from consideration of hydrogen plasma [47], a maximum in $R^2 g_{gg}(R)$ signals population of a bound state. For comparison, the quark-antiquark correlation function, i.e. $R^2 g_{q\bar{q}}(R)$, is also presented in Fig. 5. It demonstrates that there are no bound meson-like states. We can only speak about weak meson-like clustering at lower temperatures, see short-distance $q\bar{q}$ correlation at $T=193$ MeV in Fig. 4. Possible existence of medium-modified meson-like bound states was actively discussed some time ago, e.g., in Ref. [64] and later in Refs. [65, 66] based on results from lattice QCD calculations of spectral functions [67, 68]. Our result supports conclusion of Ref. [69] on the absence of $q\bar{q}$ bound states above the temperature of the phase transition. This finding is in contrast to our previous results on SU(2) group [30–33]. There well pronounced bound $q\bar{q}$ states were found just above the critical temperature, which however quickly dissolved with the temperature rise. This happens because the SU(3) plasma turns out to be essentially denser than the SU(2) one, which is a consequence of a stronger effective attraction between constituents. As a result, possible bound states in the SU(3) plasma just melt. To verify the relevance of all above discussed trends, a more refined color-, flavor-, spin-resolving analysis of the PDF's is necessary. This work is presently in progress.

Details of our path integral Monte-Carlo simulations have been discussed elsewhere in a number of papers and review articles, see, e.g. Refs. [47, 63] and references therein. The main idea of the simulations consists in constructing a Markov chain of configurations which differ by the quasiparticle coordinates. Additionally to the case of electrodynamic plasmas, here we randomly sample the color quasiparticle variables according to the SU(3) group Haar measure with the quadratic and cubic Casimir conditions until a full convergence is achieved. Another important difference from the case of the electrodynamic plasmas consists in using the relativistic measure in Feynman-Wiener path integrals associated with relativistic kinetic energy operator instead of the conventional Gaussian one arising from the non-relativistic operator. For simulations we use a cubic simulation box with periodic boundary conditions.

At the first step a dominant, i.e. maximal, $\{N\}$ -term in the sum of Eq. (6) was determined by calculations in grand canonical ensemble. This term is indeed dominant in the thermodynamic limit of the box volume $\rightarrow \infty$. Then all the calculation was performed for a canonical ensemble with thus determined $\{N\}$. In practical calculations we always take the number of particles $N = N_q + N_{\bar{q}} + N_g = 126$ and adjust the above determined proper densities of species by varying the total volume V of the box. The number of beads $n = 20$ for each particle was used. On one hand, errors in MC calculations of thermodynamic quantities related to the finite number of particle in a system with periodic boundary conditions are of order $1/N$ [47]. On other hand too large number of particle presented by

large number of beads requires large computer resources. Our choice of particle and beads numbers is a compromise between acceptable accuracy and available computer resources. Test variation of the number of beads from 15 up to 50 practically does not change results.

B. Transport coefficients

An important aspect of the strongly coupled QGP is its transport properties which strongly differ from those we would expect for weakly coupled plasmas. We use the developed approach based on Wigner formulation of quantum mechanics to calculate the QGP transport properties at strong coupling. In particular, we calculate the QGP self-diffusion constant and shear viscosity, as these quantities can be compared to respective values deduced from analysis of experimental data heavy-ion collisions and also predictions of the lattice QCD computations. More precisely, summary of shear viscosity deduced from analysis of experimental elliptic flow is presented in Ref. [73], in Ref. [74] an extensive review of theoretical works on viscosity is done, while the heavy-quark diffusion constant is available from the experiment analysis [71, 72] and QCD lattice computations [59, 70]. We anticipate that the self-diffusion and heavy-quark diffusion constants are comparable by the order of magnitude.

A natural way to obtain these transport coefficients is use of the quantum Green-Kubo relations. These relations give the transport coefficients in terms of integrals of equilibrium time-depended correlation functions. According to the Eq. (38) a self-diffusion constant D is the integral of the velocity autocorrelation function

$$\begin{aligned} D &= \lim_{t \rightarrow \infty} D(t), \\ D(t) &= \frac{1}{3N} \int_0^t d\tau \langle v(0) \cdot v(\tau) \rangle \\ \langle v(0) \cdot v(\tau) \rangle &= (2\pi)^{-6N} \int d\vec{p} d\vec{q} d\vec{\mu} d\vec{\nu} W(\vec{p} \vec{q}; \vec{\mu} \vec{\nu}; \tau; \beta) v(\vec{p}) \cdot v(\vec{q}), \end{aligned} \quad (70)$$

where the product of 3-velocities is

$$v(\vec{p}) \cdot v(\vec{q}) = \sum_{i=1}^N \frac{\vec{p}_i \cdot \vec{q}_i}{\sqrt{\vec{p}_i^2 + m_i^2} \sqrt{\vec{q}_i^2 + m_i^2}}. \quad (71)$$

Figure 6 shows examples of the velocity-velocity autocorrelation and its antiderivative function. The self-diffusion constant is a limiting value of the related antiderivative function at $t \rightarrow \infty$.




FIG. 6: Velocity autocorrelation function (left panel) and temperature-scaled self-diffusion function $T D(t)$ (right panel) versus time for two temperatures.

Analogously, the Green-Kubo relation for the shear viscosity is the integral of the autocorrelation function of the stress-energy tensor

$$\begin{aligned} \eta &= \lim_{t \rightarrow \infty} \eta(t), \\ \eta(t) &= \frac{1}{VT} \int_0^t d\tau \langle \sigma_{xy}(0) \sigma_{xy}(\tau) \rangle \end{aligned}$$

$$\langle \sigma_{xy}(0) \sigma_{xy}(\tau) \rangle = (2\pi)^{-6N} \int \overline{dpq\mu Q} \widetilde{dpq\mu Q} W(\overline{pqQ}; \widetilde{pqQ}; \tau; \beta) \sigma_{xy}(\overline{pqQ}) \sigma_{xy}(\widetilde{pqQ}), \quad (72)$$

where the off-diagonal stress-energy tensor is

$$\sigma_{xy}(pqQ) = \sum_{i=1}^N \frac{p_{i,x} p_{i,y}}{\sqrt{\mathbf{p}_i^2 + m_i^2}} - \frac{1}{2} \sum_{i \neq j}^N q_{ij,x} \frac{\partial U(qQ)}{\partial q_{ij,y}}, \quad (73)$$

here $\mathbf{q}_{ij} = \mathbf{q}_i - \mathbf{q}_j$, and U is the sum of the color Kelbg potentials defined by Eqs. (62) and (63) with $n = 0$. Examples of the stress-energy-tensor autocorrelation and its antiderivative functions are presented in Figure 7. The shear viscosity is defined by limiting value of the related antiderivative function at $t \rightarrow \infty$.




FIG. 7: Temperature-scaled stress-energy-tensor autocorrelation function (left panel) and shear-viscosity function $\eta(t)/T^3$ (right panel) versus time for two temperatures.




FIG. 8: Left panel: The self-diffusion constant [self-diffusion, Wigner dynamics] as a function of temperature, compared with heavy-quark diffusion constants predicted by QCD lattice computations [70] [c-quark, lattice (Ding et al. 2011)] and [59] [c-quark, lattice (Banerjee et al. 2012)], and deduced from analysis of experimental data [71] [c-quark, “exp.” (Gossiaux et al. 2012)] and [72] [c-quark, “exp.” (He et al. 2012)].

Right panel: The ratio of shear viscosity to entropy density as a function of temperature [shaded area marked “Wigner dynamics”]. Horizontal dashed lines indicate the range of constraint on the viscosity-to-entropy ratio deduced from numerous hydrodynamical simulations of heavy-ion experimental data, as summarized in Ref. [73].

The self-diffusion constant and the viscosity-to-entropy ratio are presented in Fig. 8 as a function of temperature. The entropy density s is taken from results of our PIMC calculations presented in Fig. 2. Our results are presented in

the form of bands. The width of the bands represents the theoretical uncertainty associated with the oscillations of the antiderivative functions at large times, see Figs. 6 and 7. Slowly decaying oscillations of the time correlation functions are inherent in liquid-like systems of strongly interacting particles in contrast to exponentially decaying oscillations in gas-like systems. As known from hydrogen plasma, these oscillations arise because of quasiclosed chaotic orbits and are caused by strong interparticle interaction. In liquids these oscillations decay according to a power law rather than exponentially. Therefore, extremely long (in time) trajectories are required for more accurate calculations of the diffusion constant and viscosity. Due to CPU time limitations, we had to stop our computations before the decay of these oscillations was completed.

Unfortunately, the self-diffusion constant is unavailable from other calculations. Therefore, we compare it with heavy-quark diffusion constant, anticipating that these are of the same order of magnitude. The heavy-quark diffusion constant is available from recent QCD lattice computations [59, 70], which are presented in the left panel of Figs. 8. Our results (labeled as “Wigner dynamics”) well agree with lattice data of Ref. [59], while essentially overestimate those of Ref. [70]. The heavy-quark diffusion constant is also available from analysis of experiments on heavy-quark quenching in ultrarelativistic heavy ions collisions at RHIC. For comparison we took two recent works on the subject [71, 72]. Here the results are also rather diverse. Estimates of Ref. [72] are well comparable with our result and lattice data of Ref. [59], while estimate of Ref. [71, 72] is considerably lower and better conforms to lattice results of Ref. [70].

Our results on the shear viscosity are presented in the right panel of Figs. 8. As seen, the obtained values of viscosity are in the range of those deduced from the analysis of experimental elliptic flow in ultrarelativistic heavy ions collisions at RHIC, as summarized in Ref. [73]. Lattice data on the shear viscosity in the realistic case of the SU(3) group are not available, an extensive review of theoretical works on viscosity within QCD-motivated models is done in Ref. [74]. As seen, the minimum of the viscosity-to-entropy ratio is reached at a temperature above the expected phase transition rather than in the phase transition range, as is commonly expected. This minimum turns out to be quite shallow. The value of the viscosity-to-entropy ratio at the minimum is very close (from above) to the lower bound of $\eta/S = 1/4\pi$ for this quantity [75], often referred to as the KSS bound. With the temperature decrease, i.e. towards the hadronic phase, the viscosity rapidly rises.

V. CONCLUSION

In this paper we demonstrated that color quantum Monte-Carlo (PIMC) simulations based on the quasiparticle model of the QGP are able to reproduce the lattice equation of state at zero baryon chemical potential at realistic model parameters (i.e. quasiparticle masses and coupling constant) even near the critical temperature and also yields valuable insight into the internal structure of the QGP. In our simulations we have introduced a new relativistic path integral measure and have developed a procedure of sampling color quasiparticle variables according to the SU(3) group Haar measure with appropriate Casimir conditions. Unfortunately, convergence of our calculations becomes poor in the range of the expected phase transition because the scheme suffers from jumps between stable and metastable states which turn out to be almost equally probable in this range. Our results indicate that the QGP reveals *quantum liquid-like* (rather than gas-like) properties up to the highest considered temperature of 525 MeV.

Short-distance correlations in the computed pair distribution functions of gluon-gluon and gluon-(anti)quark pairs displays the formation of clusters. In case of the gluon-gluon clusters we can even speak about gluon-gluon bound states, i.e. glueballs, at temperatures just above the phase transition. The possible existence of medium-modified meson-like bound states was actively discussed some time ago [65, 66]. Our result supports the conclusion of Ref. [69] on the absence of $q\bar{q}$ bound states above the temperature of the phase transition. This finding is in contrast to our previous results on the SU(2) group [30–33]. There well pronounced bound $q\bar{q}$ states were found just above the critical temperature, which however quickly dissolved with the temperature rise. This happens because the SU(3) plasma turns out to be essentially denser than the SU(2) one, which is a consequence of a stronger effective attraction between the constituents. As a result, possible meson-like bound states in the SU(3) plasma just melt.

The PIMC method is not able to yield transport properties of the QGP. A way to access these is to develop a classical color molecular dynamics simulation [15], where quantum effects are included phenomenologically via a short-range potential. In contrast to these classical MD simulations [15], we have developed a more rigorous approach based on the combination of Feynman and Wigner formulations of quantum dynamics. The basic ideas of this approach have been briefly reported in Ref. [34]. In this paper we gave a more detailed description. In particular, this approach allowed us to calculate the self-diffusion coefficient and the viscosity of the strongly coupled QGP. Since the self-diffusion constant is unavailable from other calculations, we compared it with the heavy-quark diffusion constant, anticipating that these are of the same order of magnitude. The heavy-quark diffusion constant is available from recent QCD lattice computations [59, 70] and also from an analysis of the heavy-quark quenching in experiments on ultrarelativistic heavy ions collisions at RHIC. For comparison we took two recent works on such an analysis [71, 72].

Unfortunately the above mentioned lattice and heavy-quark-quenching results are rather diverse. Our self-diffusion constant well agrees with lattice data of Ref. [59] and estimates of Ref. [72], while essentially overestimates those of Refs. [70] and [70].

Our results on the shear viscosity are in the range of those deduced from the analysis of the experimental elliptic flow in ultrarelativistic heavy ions collisions at RHIC, as summarized in Ref. [73], i.e. in terms of the viscosity-to-entropy ratio, $1/4\pi \lesssim \eta/S < 2.5/4\pi$, in the temperature range from 170 to 440 MeV. The minimum of the viscosity-to-entropy ratio is reached at a temperature (≈ 300 MeV), above the expected phase transition rather than in the phase transition range as commonly expected. This minimum turns out to be very shallow. The value of the viscosity-to-entropy ratio at the minimum is very close (from above) to the lower bound of $\eta/S = 1/4\pi$ for this quantity [75], i.e. to the KSS bound. With the temperature decrease, i.e. towards the hadronic phase, the viscosity rapidly rises.

Our present analysis is still confined only to the case of zero baryon chemical potential. Simulations at nonzero baryon chemical potentials are in progress.

We acknowledge stimulating discussions with P. Levai, D. Blaschke, R. Bock, H. Stoecker, and D.N.Voskresensky. Y.B.I. was partially supported by grant of the Russian Ministry of Science and Education NS-215.2012.2.

Appendix I: Integration over SU(3) group Haar measure

In this appendix we explain details of integration over SU(3) Haar measure $d\mu Q$ in Eq. (6). The measure for single color charge in case of the SU(3) group is [10, 76–78]

$$d\mu Q = d^8 Q \delta(Q_a Q_a - \check{q}_2) \delta(d_{abc} Q_a Q_b Q_c - \check{q}_3) \quad (74)$$

with summation over $a, b, c = 1, \dots, 8$ and constants d_{abc} given in Table 1. For the SU(N) group the quadratic \check{q}_2 and cubic \check{q}_3 Casimirs are $\check{q}_2 = (N^2 - 1)C^2$ with $C_2 = N$ for gluons and $C_2 = 1/2$ for quarks and antiquarks, $\check{q}_3 = 0$ for gluons and $\check{q}_3 = (N^2 - 4)(N^2 - 1)/4$ for quarks, for antiquarks \check{q}_3 has opposite sign. In fact, the normalization constant c_R depends on \check{q}_2 and \check{q}_3 Casimirs.

For random sampling of the Q variable in Monte-Carlo integration in Eq. (6) we change to the related canonical Darboux variables for the SU(3) group. The set of the canonical variables $[\phi\pi] = [(\phi_\alpha, \pi_\alpha, \alpha = 1, 2, 3)]$ is defined by the canonical Poisson bracket

$$\{A, B\}_{PB} = \frac{\partial A}{\partial r} \frac{\partial B}{\partial p} - \frac{\partial A}{\partial p} \frac{\partial B}{\partial r} + \frac{\partial A}{\partial \phi} \frac{\partial B}{\partial \pi} - \frac{\partial A}{\partial \pi} \frac{\partial B}{\partial \phi} \quad (75)$$

where r and p are conventional coordinate and momentum, respectively, and obey

$$\{r_\alpha, p_\gamma\}_{PB} = \delta_{\alpha\gamma} \quad \{\phi_\alpha, \pi_\gamma\}_{PB} = \delta_{\alpha\gamma} \quad (76)$$

The color variables Q_a form a representation of SU(3). In terms canonical variables their Poisson bracket reads

$$\{Q_a, Q_b\}_{PB} = \check{f}_{abc} Q_c \quad (77)$$

where \check{f}_{abc} are the structure constants of SU(3) given in Table 1.

The explicit transformations to canonical variables are given [10, 76–78] by expressions:

$$\begin{aligned} Q_1 &= \pi_+ \pi_- \cos \phi_1, & Q_2 &= \pi_+ \pi_- \sin \phi_1, & Q_3 &= \pi_1, \\ Q_4 &= C_{++} \pi_+ A + C_{+-} \pi_- B, & Q_5 &= S_{++} \pi_+ A + S_{+-} \pi_- B, \\ Q_6 &= C_{-+} \pi_- A + C_{--} \pi_+ B, & Q_7 &= S_{-+} \pi_- A + S_{--} \pi_+ B, & Q_8 &= \pi_2 \end{aligned} \quad (78)$$

in which we have used definitions:

$$\begin{aligned} \pi_+ &= \sqrt{\pi_3 + \pi_1} & \pi_- &= \sqrt{\pi_3 - \pi_1} \\ C_{\pm\pm} &= \cos \left[\frac{1}{2} \left(\pm \phi_1 + \sqrt{3} \phi_2 \pm \phi_3 \right) \right] & S_{\pm\pm} &= \sin \left[\frac{1}{2} \left(\pm \phi_1 + \sqrt{3} \phi_2 \pm \phi_3 \right) \right] \end{aligned} \quad (79)$$

and A and B are given by

$$\begin{aligned} A &= \frac{1}{2\pi_3} \sqrt{\left(\frac{J_1 - J_2}{3} + \pi_3 + \frac{\pi_2}{\sqrt{3}} \right) \left(\frac{J_1 + 2J_2}{3} + \pi_3 + \frac{\pi_2}{\sqrt{3}} \right) \left(\frac{2J_1 + J_2}{3} - \pi_3 - \frac{\pi_2}{\sqrt{3}} \right)} \\ B &= \frac{1}{2\pi_3} \sqrt{\left(\frac{J_2 - J_1}{3} + \pi_3 - \frac{\pi_2}{\sqrt{3}} \right) \left(\frac{J_1 + 2J_2}{3} - \pi_3 + \frac{\pi_2}{\sqrt{3}} \right) \left(\frac{2J_1 + J_2}{3} + \pi_3 - \frac{\pi_2}{\sqrt{3}} \right)} \end{aligned} \quad (80)$$

In this expression the set Q_1, Q_2, Q_3 forms an $SU(2)$ subgroup with quadratic Casimir $Q_1^2 + Q_2^2 + Q_3^2 = \pi_3^2$. Let us note that two Casimirs depend only on J_1 and J_2 . They can be computed using the values given in Table 1 as

$$\begin{aligned} Q_a Q_a &= \frac{1}{3} (J_1^2 + J_1 J_2 + J_2^2) \\ d_{abc} Q_a Q_b Q_c &= \frac{1}{18} (J_1 - J_2) (J_1 + 2J_2) (2J_1 + J_2) \end{aligned} \quad (81)$$

The phase space color measure for $SU(3)$, given in Eq. (74) can be transformed to the new coordinates through use of Eq. (78) and evaluation of the Jacobian

$$\left| \frac{\partial Q}{\partial(\phi, \pi)} \right| = \frac{\sqrt{3}}{48} J_1 J_2 (J_1 + J_2) \quad (82)$$

Then the measure reads

$$\begin{aligned} d\mu Q &= d\phi_1 d\phi_2 d\phi_3 d\pi_1 d\pi_2 d\pi_3 dJ_1 dJ_2 \frac{\sqrt{3}}{48} J_1 J_2 (J_1 + J_2) \\ &\times \delta \left(\frac{1}{3} ((J_1)^2 + J_1 J_2 + (J_2)^2) - \check{q}_2 \right) \delta \left(\frac{1}{18} (J_1 - J_2) (J_1 + 2J_2) (2J_1 + J_2) - \check{q}_3 \right) \end{aligned} \quad (83)$$

Since the two Casimirs are independent, the δ -functions fix both J^1 and J^2 . After integration over J^1 and J^2 the Eq. (83) gives a proper canonical volume element $d\phi d\pi$. Thus applying Metropolis algorithm to $\phi\pi$ variables we can construct Markovian chain in $\phi\pi$ phase space and obtain random color variables Q for calculation partition function according to the $SU(3)$ group Haar measure with two Casimir conditions.

TABLE I: Non-zero \check{f}_{abc} and d_{abc} constants of the group $SU(3)$

		\check{f}_{abc}	\check{f}_{123}	\check{f}_{147}	\check{f}_{156}	\check{f}_{246}	\check{f}_{257}	\check{f}_{345}	\check{f}_{367}	\check{f}_{458}	\check{f}_{678}						
		values	1	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$						
d_{abc}	d_{118}	d_{146}	d_{157}	d_{228}	d_{247}	d_{256}	d_{338}	d_{344}	d_{355}	d_{366}	d_{377}	d_{448}	d_{558}	d_{668}	d_{778}	d_{888}	
values	$\frac{1}{\sqrt{3}}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{\sqrt{3}}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{\sqrt{3}}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{1}{\sqrt{3}}$	

Appendix II: Lebesgue-Dirac delta theorem

Let f be a summable function of real argument such that $\int f(x)dx = I_0$. Then $Mf(M(x - x')) \rightarrow I_0\delta(x - x')$ when $M \rightarrow \infty$.

Proof: Let φ be any test function. Then

$$\begin{aligned} \lim_{M \rightarrow \infty} \int Mf(M(x - x')) \varphi(x) dx &= \lim_{M \rightarrow \infty} \int f(s) \varphi(s/M + x') ds \\ &= \int f(s) \left(\lim_{M \rightarrow \infty} \varphi(s/M + x') \right) ds = \varphi(x') \int f(s) ds = I_0 \varphi(x') \end{aligned}$$

-
- [1] E. Shuryak, Prog. Part. Nucl. Phys. **62**, 48 (2009) [arXiv:0807.3033 [hep-ph]]; *ibid.* **53**, 273 (2004) [arXiv:hep-ph/0312227].
 - [2] J. Schukraft, arXiv:1112.0550 [arXiv:hep-ex].
 - [3] C. Shen and U. Heinz, Phys. Rev. C **85**, 054902 (2012) [arXiv:1202.6620 [nucl-th]].
 - [4] Z. Fodor and S. D. Katz, arXiv:0908.3341 [hep-ph].
 - [5] S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S. D. Katz, S. Krieg, C. Ratti, K. K. Szabo, JHEP **1011**, 077 (2010) [arXiv:1007.2580 [hep-lat]].
 - [6] M. Cheng *et al.*, Phys. Rev. D **81**, 054504 (2010) [arXiv:0911.2215 [hep-lat]].
 - [7] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, Nature **443**, 675 (2006) [hep-lat/0611014].
 - [8] H. B. Meyer, Phys. Rev. D **76**, 101701 (2007) [arXiv:0704.1801 [hep-lat]].
 - [9] P. Hartmann, Z. Donko, P. Levai, G.J. Kalman, J. Phys. **A42**, 214029 (2006); Nucl. Phys. **A774**, 881-884, (2006) [arXiv:nucl-th/0601017].

- [10] D. F. Litim and C. Manuel, Phys. Rev. Lett. **82**, 4981 (1999) [arXiv:hep-ph/9902430]; Nucl. Phys. B **562**, 237 (1999) [arXiv:hep-ph/9906210]; Phys. Rev. D **61**, 125004 (2000) [arXiv:hep-ph/9910348]; Phys. Rep. **364**, 451 (2002) [arXiv:hep-ph/0110104].
- [11] M. Laine and C. Manuel, Phys. Rev. D **65**, 077902 (2002) [arXiv:hep-ph/0111113].
- [12] C. Manuel and S. Mrowczynski, Phys. Rev. D **68**, 094010 (2003) [arXiv:hep-ph/0306209].
- [13] P. F. Kelly, Q. Liu, C. Lucchesi and C. Manuel, Phys. Rev. D **50**, 4209 (1994) [arXiv:hep-ph/9406285].
- [14] M. Hofmann, M. Bleicher, S. Scherer, *et al.*, Phys. Lett. B **478**, 161 (2000) [arXiv:nucl-th/9908030].
- [15] B.A. Gelman, E. V. Shuryak, and I. Zahed, Phys. Rev. C **74**, 044908 (2006) [nucl-th/0601029]; *ibid.* **74**, 044909 (2006) [arXiv:nucl-th/0605046].
- [16] S. Cho and I. Zahed, Phys. Rev. C **79**, 044911 (2009) [arXiv:0812.1736 [nucl-th]]; *ibid.* **80**, 014906 (2009) [arXiv:0812.1741 [nucl-th]]; *ibid.* **82**, 054907 (2010) [arXiv:0910.2666 [nucl-th]]; *ibid.* **82**, 064904 (2010) [arXiv:0910.1548 [nucl-th]]; *ibid.* **82**, 044905 (2010) [arXiv:0909.4725 [nucl-th]]; K. Dusling and I. Zahed, Nucl. Phys. A **833**, 172 (2010) [arXiv:0904.0169 [nucl-th]].
- [17] M.H. Thoma, IEEE Trans. Plasma Science **32**, 738 (2004).
- [18] A. Filinov, M. Bonitz, and W. Ebeling, J. Phys. A **36**, 5957 (2003).
- [19] G. Kelbg, Ann. Phys. (Leipzig) **12**, 219 (1962); **13**, 354 (1963).
- [20] K. Dusling and C. Young, arXiv:0707.2068 [nucl-th].
- [21] A. Filinov, V. Golubnychiy, M. Bonitz, *et al.*, Phys. Rev. E **70**, 046411 (2004)
- [22] M. Bonitz, A. Filinov, V. Golubnychiy, *et al.*, J. Phys. A **36**, 5921 (2003); Phys. **15**, 055704 (2008).
- [23] V. S. Filinov, M. Bonitz, W. Ebeling, and V. E. Fortov, Plasma Phys. Control. Fusion **43**, 743 (2001).
- [24] V. S. Filinov, M. Bonitz, and V. E. Fortov, JETP Lett. **72**, 245 (2000).
- [25] M. Bonitz, V. S. Filinov, V. E. Fortov, *et al.*, Phys. Rev. Lett. **95**, 235006 (2005).
- [26] V. S. Filinov, M. Bonitz, P. R. Levashov, *et al.*, J. Phys. A **36**, 6069 (2003).
- [27] M. Bonitz, V. S. Filinov, V. E. Fortov, *et al.*, J. Phys. A **39**, 4717 (2006).
- [28] V. S. Filinov, H. Fehske, M. Bonitz, *et al.*, Phys. Rev. E **75**, 036401 (2007).
- [29] V. S. Filinov, V. E. Fortov, P. R. Levashov, Y. B. Ivanov, M. Bonitz, V. E. Fortov and P. R. Levashov, Phys. Lett. A **376**, 1096 (2012) [arXiv:1203.2191 [hep-ph]].
- [30] V. S. Filinov, M. Bonitz, Y. B. Ivanov, V. V. Skokov, P. R. Levashov and V. E. Fortov, Contrib. Plasma Phys. **52**, 135 (2012).
- [31] V. S. Filinov, Y. B. Ivanov, M. Bonitz, P. R. Levashov and V. E. Fortov, Phys. Atom. Nucl. **74**, 1364 (2011) [arXiv:1101.2089 [nucl-th]].
- [32] V. S. Filinov, Y. B. Ivanov, M. Bonitz, P. R. Levashov and V. E. Fortov, Phys. Atom. Nucl. **75**, 693 (2012).
- [33] V. S. Filinov, M. Bonitz, Y. B. Ivanov, V. V. Skokov, P. R. Levashov and V. E. Fortov, Contrib. Plasma Phys. **49**, 536 (2009) [arXiv:0905.0324 [nucl-th]].
- [34] V. S. Filinov, M. Bonitz, Y.B. Ivanov, V.V. Skokov, P.R. Levashov, and V.E. Fortov, Contrib. Plasma. Phys. **51**, N4, 322-327 (2011).
- [35] P. Petreczky, F. Karsch, E. Laermann, S. Stickan, and I. Wetzorke, Nucl. Phys. Proc. Suppl. **106**, 513 (2002) [arXiv:hep-lat/0110111].
- [36] J. Liao and E. V. Shuryak, Phys. Rev. D **73**, 014509 (2006).
- [37] F. Karsch and M. Kitazawa, Phys. Rev. D **80**, 056001 (2009) [arXiv:0906.3941 [hep-lat]].
- [38] S. K. Wong, Nuovo Cimento A **65**, 689 (1970).
- [39] R. D. Pisarski, Phys. Rev. Lett. **63**, 1129 (1989).
- [40] E. Braaten and R. D. Pisarski, Nucl. Phys. **B337**, 569 (1990); **B339**, 310 (1990).
- [41] J. Blaizot and E. Iancu, Phys. Rept. **359**, 355 (2002) [arXiv:hep-ph/0101103].
- [42] E. E. Salpeter and H. A. Bethe, Phys. Rev. **84**, 1232, (1951).
- [43] E. E. Salpeter, Phys. Rev. **87**, 328, (1952).
- [44] W. Lucha, F. F. Schoberl, and D. Gromes, Phys. Rep. **200**, 127, (1991).
- [45] W. Lucha, F. F. Schoberl, Int. J. Mod. Phys. **A7**, 6431, (1992).
- [46] R. P. Feynman, and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- [47] V. M. Zamalin, G. E. Norman, and V. S. Filinov, *The Monte-Carlo Method in Statistical Thermodynamics* (Nauka, Moscow, 1977), (in Russian).
- [48] V. Tatarskii, Sov. Phys. Uspekhi **26**, 311 (1983).
- [49] V. Filinov, Y. Medvedev, V. Kamskii, Mol. Phys. **85**(4), 711 (1995).
- [50] G. Ciccotti, C. Pierleoni, F. Capuani, V. Filinov, Comp. Phys. Comm. **121-122**, 452 (1999).
- [51] S. Cho and I. Zahed, Phys. Rev. C **82**, 054907 (2010) [arXiv:0910.2666 [nucl-th]].
- [52] D.N. Zubarev, *Nonequilibrium Statistical Thermodynamics*, Plenum Press, New York/London 1974.
- [53] J.D. Doll, D.L. Freeman, T.L. Beck, Adv. Chem. Phys. **78**, 61-127 (1990).
- [54] V.S. Filinov, J. Mol. Phys. **88**, 1517, 1529 (1996).
- [55] V. Filinov, P. Thomas, I. Varga, T. Meier, M. Bonitz, V. Fortov and S. Koch, Phys. Rev. **B65**, 165124, (2002).
- [56] D. V. Shirkov and I. L. Solovtsov, Phys. Rev. Lett. **79** (1997) 1209; Nucl. Phys. Proc. Suppl. **64**, 106 (1998).
- [57] G. M. Prospero, M. Raciti, and C. Simolo, Prog. Part. Nucl. Phys. **58**, 387 (2007) [arXiv:hep-ph/0607209].
- [58] O. Kaczmarek, F. Karsch, F. Zantow, and P. Petreczky, Phys. Rev. D **72**, 059903 (2005) [arXiv:hep-lat/0406036].
- [59] D. Banerjee, S. Datta, R. Gavai and P. Majumdar, Phys. Rev. D **85**, 014510 (2012) [arXiv:1109.5738 [hep-lat]].
- [60] A. Nakamura, T. Saito and S. Sakai, Phys. Rev. D **69**, 014506 (2004) [arXiv:hep-lat/0311024].

- [61] A. Peshier, B. Kampfer, and O.P. Pavlenko, Phys. Rev. D **54**, 2399 (1996).
- [62] Yu. B. Ivanov, V. V. Skokov, and V. D. Toneev, Phys. Rev. D **71**, 014005 (2005). [arXiv:hep-ph/0410127].
- [63] A. V. Filinov, V. S. Filinov, Yu. E. Lozovik and M. Bonitz, *Introduction to Computational Methods for Many-Body Physics*, Ed. by M. Bonitz and D. Semkat (Rinton Press, Princeton, 2006).
- [64] V. I. Yukalov and E. P. Yukalova, Physica A **243**, 382 (1997); Phys. Part. Nucl. 28, 37 (1997).
- [65] E. V. Shuryak and I. Zahed, Phys. Rev. C **70**, 021901 (2004) [arXiv:hep-ph/0307267]; Phys. Rev. D **70**, 054507 (2004) [arXiv:hep-ph/0403127].
- [66] G. E. Brown, B. A. Gelman, and M. Rho, arXiv:nucl-th/0505037.
- [67] M. Asakawa, T. Hatsuda and Y. Nakahara, Prog. Part. Nucl. Phys. **46**, 459 (2001) [arXiv:hep-lat/0011040]; Nucl. Phys. A **715**, 863 (2003) [Nucl. Phys. Proc. Suppl. **119**, 481 (2003)] [arXiv:hep-lat/0208059]; M. Asakawa and T. Hatsuda, Phys. Rev. Lett. **92**, 012001 (2004) [arXiv:hep-lat/0308034].
- [68] S. Datta, F. Karsch, P. Petreczky and I. Wetzorke, Phys. Rev. D **69**, 094507 (2004) [arXiv:hep-lat/0312037].
- [69] V. Koch, A. Majumder, and J. Randrup, Phys. Rev. Lett. **95**, 182301 (2005) [arXiv:nucl-th/0505052].
- [70] H. T. Ding, A. Francis, O. Kaczmarek, F. Karsch, H. Satz and W. Soeldner, Phys. Rev. D **86**, 014509 (2012) [arXiv:1204.4945 [hep-lat]].
- [71] P. Gossiaux, J. Aichelin and T. Gousset, Prog. Theor. Phys. Suppl. **193**, 110 (2012) [arXiv:1201.4038 [hep-ph]]; P. B. Gossiaux, J. Aichelin, M. Bluhm, T. Gousset, M. Nahrgang, S. Vogel and K. Werner, arXiv:1207.5445 [hep-ph].
- [72] M. He, R. J. Fries and R. Rapp, arXiv:1204.4442 [nucl-th].
- [73] S. A. Bass, Nucl. Phys. A **862-863**, 174 (2011).
- [74] A. S. Khvorostukhin, V. D. Toneev and D. N. Voskresensky, Nucl. Phys. A **845**, 106 (2010) [arXiv:1003.3531 [nucl-th]].
- [75] P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005) [arXiv:hep-th/0405231].
- [76] K. Johnson, Ann. Phys. **192**, 104, (1989).
- [77] P.F. Kelly, Q. Liu, C. Lucchesi and C. Manuel, Phys. Rev. Lett. **72**, 3461, (1994).
- [78] P.F. Kelly, Q. Liu, C. Lucchesi and C. Manuel, Phys. Rev. **D50**, 4209, (1994).